

### GENERAL INFORMATION

Du Pont "Trysben" 200 Weed Killer is a liquid to be diluted with water for use as a spray for non-selective control of certain undesirable broadleaf weeds and certain species of woody plants. Both foliage and root absorption contribute to plant kill. "Trysben" 200 is intended for use on utility, highway, pipeline, railroad right of ways, and other industrial sites; ditchbanks; fencerows; and around farm buildings. Application as directed will result in loss of soil productivity for an extended period.

### DIRECTIONS

"Trysben" 200 should be used only in accordance with recommendations on this label, or in separate Du Pont Agricultural Bulletins available through local dealers.

### HOW TO APPLY

"Trysben" 200 should be applied in water as a coarse spray in ground equipment using low or moderate pressure. If the spray is mixed thoroughly after dilution to provide a uniform solution, equipment without agitation may be used. Where ground cover is light, use at least 50 gals. of spray solution per acre. If ground cover is heavy, a volume of 100 or more gals. per acre is recommended. Spray volume should be sufficient to provide uniform and thorough coverage of foliage and soil.

### WHEN TO APPLY

Best results are obtained if application is made after foliage has appeared in the spring, but before growth has progressed sufficiently to retain more than half of the spray. Earlier or later applications also are effective, but rates near the upper limit of the recommended dosage range may be required. Applications should be made preferably during periods when moderate rainfall can be expected.

### NOXIOUS PERENNIAL WEEDS

To control certain deep-rooted perennial weeds such as field bindweed, leafy spurge, Canada thistle, Russian knapweed, bur ragweed, blueweed, and

UNDER THE FEDERAL INSECTICIDE,  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
ED UNDER NO.



REG. U. S. PAT. OFF.

# TRYSBEN<sup>®</sup> 200

## WEED KILLER

CONTAINS 2 POUNDS TRICHLORO BENZOIC ACID EQUIVALENT PER GALLON

**ACTIVE INGREDIENTS**..... 26.1%  
 Dimethylamine salt of 2,3,6-trichlorobenzoic acid. 15.7%  
 Dimethylamine salts of  
 other chlorinated benzoic acids..... 10.4%  
 (Total trichlorobenzoic acid equivalent..... 21.7%)  
**INERT INGREDIENTS**..... 73.9%

U.S. Pats. 2,848,470 & 3,081,162  
 USDA Reg. No. 352-250

Keep out of reach of children.

**CAUTION! MAY IRRITATE SKIN AND EYES.**

Avoid contact with skin, eyes, and clothing.

In case of contact, flush with plenty of water;  
 for eyes, get medical attention.

IF EXPOSED TO PROLONGED COLD, PLACE IN WARM  
 STORAGE (50 TO 80° F.) FOR SEVERAL HOURS AND AGITATE BEFORE USING.

## 30 U.S. GALLONS

**IMPORTANT**—Injury to or loss of desirable trees or other plants may result from failure to observe the following: Do not apply, or drain or flush equipment, on or near desirable trees or other plants, or in areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, tennis courts, or similar areas. Do not use area actually treated for crop production. Spray drift, and possibly vapors from treated areas, may injure susceptible crops such as beans of all types, tomatoes, peas, cotton, tobacco, various ornamental plants, orchard and vine crops. Do not apply to areas where nearby fields are to be used for growing susceptible crops during the same season. Do not contaminate domestic or irrigation waters. Keep from contact with fertilizers, insecticides, fungicides, and seeds. Thoroughly clean application equipment after use. Do not use equipment with which "Trysben" 200 has been applied for spraying insecticides, fungicides, selective herbicides, and fertilizers as plant injury may result.

Do not re-use container. Return to qualified drum reconitioner, or crush and bury when empty.

climbing milkweed, apply 5 to 10 gals. "Trysben" 200 (10 to 20 lbs. acid equivalent) per acre. For small areas, use 1 to 2 pts. per 1,000 sq. ft. For round weed patches, mix "Trysben" 200 at rate of ½ to 1 gal. per 15 gals. of water and apply spray mixture at following rate:

DISTANCE ACROSS AREA TO BE TREATED (weed patch plus 10 feet each side)	AMOUNT OF SPRAY MIXTURE
25 Feet	2 Gallons
50 Feet	7 Gallons
75 Feet	15 Gallons
100 Feet	27 Gallons

### MIXED BROADLEAF WEEDS

To control a wide range of perennial and annual broadleaf weeds for one year or longer, apply 10 to 15 gals. "Trysben" 200 (20 to 30 lbs. acid equivalent) per acre. Weeds controlled include the above noxious perennials and others such as dandelion, plantain, dock, dog fennel, ragweed, and wild lettuce. For small areas, use 2 to 3 pts. per 1,000 sq. ft.

### WOODY VINES

To control certain woody vines such as trumpet vine, honeysuckle, kudzu, and smilax, apply 5 to 10 gals. "Trysben" 200 (10 to 20 lbs. acid equivalent) per acre.

### WOODY BRUSH

To control certain species such as conifers (spruce, balsam fir, cedar, and pine), wild roses (including Macartney or Cherokee rose), sumac, persimmon, sassafras, black locust, and hackberry, use 2 gals. "Trysben" 200 (4 lbs. acid equivalent) per 100 gals. of water. Thoroughly spray foliage and stems, as well as the ground under the plants.

**NOTICE TO BUYER**—Seller warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. This warranty does not extend to use of this product contrary to label use directions, or under abnormal use conditions, or under conditions not reasonably foreseeable to seller; buyer assumes all risk of any such use. Seller makes no other warranties, express or implied.

MADE IN U.S.A.

LITHO IN U.S.A.

MANUFACTURED FOR

**E. I. DU PONT DE NEMOURS & COMPANY (INC.) INDUSTRIAL AND BIOCHEMICALS DEPT., WILMINGTON, DELAWARE**