

### Preplant Incorporation

Apply the tank mix combinations of "Lorox", "Lexone" DF and "Prowl" within 7 days of planting soybeans. Incorporate the herbicides thoroughly into the top 2 inches of soil; to insure shallow incorporation of 2 inches or less, pull-type tillage equipment should be set to work the soil no deeper than 4 inches.

Plant seed 1 1/2" to 2" deep on flat or raised seedbeds only or injury to the crop may result; soil should be well prepared and as free as possible from trash and clods after planting.

### Use Rates Per Acre

<u>Soil Texture</u>	<u>1/2 to 3% Organic Matter</u>	<u>3 to 6% Organic Matter</u>
<u>Coarse</u>		
Sandy loam	1/3 to 1/2 lb. "Lorox" + 1/6 to 1/4 lb. "Lexone" DF <sup>TM</sup> + 1 to 1 1/2 pts. "Prowl"	1/3 to 3/4 lb. "Lorox" + 1/4 to 1/3 lb. "Lexone" DF <sup>TM</sup> + 1 1/2 pts. "Prowl"
<u>Medium</u>		
Silt, Sandy clay, Loam, Silt loam, Sandy clay loam	1/2 to 3/4 lb. "Lorox" + 1/4 to 1/3 lb. "Lexone" DF <sup>TM</sup> + 1 1/2 pts. "Prowl"	3/4 to 1 1/2 lbs. "Lorox" + 1/3 to 1/2 lb. "Lexone" DF <sup>TM</sup> + 1 1/2 pts. "Prowl"
<u>Fine</u>		
Clay loam, Clay, Silty clay, Silty Clay loam	3/4 to 1 1/2 lbs. "Lorox" + 1/3 to 1/2 lb. "Lexone" 4L + 1 1/2 to 2 pts. "Prowl"	1 1/2 to 2 lbs. "Lorox" + 1/2 to 2/3 lb. "Lexone" 4L + 1 1/2 to 2 pts. "Prowl"

Do not use on sand nor loamy sand with less than 1% organic matter. Do not use on any soils with less than 1/2% organic matter as crop injury may result. Do not use on muck or peat.

Do not overlap or over apply. Misapplication or improper soil incorporation can result in erratic weed control or crop injury. Seedling disease, cold weather, deep planting (more than 2"), excessive moisture, high soil pH (7.5 or higher), high salt concentration, or drought may weaken soybean seedlings and increase possibility of crop injury.



# AGRICULTURAL BULLETIN




LOROX® HERBICIDE  
CARROTS  
FLORIDA

SUPPLEMENTAL LABELING  
EPA REG. NO. 352-270

LOROX® HERBICIDE  
FOR PREEMERGENCE USE IN CARROTS  
IN THE STATE OF FLORIDA

### DIRECTIONS FOR USE



It is a violation of federal law to use this product  
in a manner inconsistent with its labeling.

Du Pont "Lorox" Herbicide is effective for use as a preemergence treatment in carrots in the state of Florida for the control of germinating weeds such as barnyardgrass (watergrass), carpetweed, chickweed, crabgrass, foxtail (including giant foxtail), goosegrass, lambsquarters, mustard, pigweed, purslane, and smartweed. Treatment will not control established perennials such as Canada thistle and purple nutsedge.

### HOW TO USE

Preemergence Application (Florida) - Apply "Lorox" after planting but before carrots emerge; plant seed at least 1/2" deep. Make a single application at the rate of 1 to 2 lbs. per acre (for band treatment, use proportionately less). Use the lower rate on lighter soils (low in clay or organic matter) and the higher rate on heavier soils (high in clay or organic matter).

Note: "Lorox" is also effective for use in carrots as postemergence application; see package label for use instructions. Carrots may be treated with multiple applications (pre- and/or postemergence) provided the total does not exceed 4 lbs. "Lorox" per acre.

### IMPORTANT

BEFORE USING "LOROX", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.