

SPECIMEN LABEL

87% OF ACTUAL SIZE

BEST AVAILABLE COPY



REGISTERED TRADEMARK

HYVAR X

HERBICIDE

WETTABLE POWDER

ACTIVE INGREDIENT:

Bromacil (5-bromo-3 sec-butyl-6-methyluracil)

80%

INERT INGREDIENTS

20%

Keep out of reach of children.
PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS

CAUTION! MAY IRRITATE EYES, NOSE, THROAT, AND SKIN

Avoid breathing dust. Avoid contact with skin, eyes or clothing.
Wash thoroughly with soap and water after handling.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

STATEMENT OF PRACTICAL TREATMENT

If in eyes: Flush with plenty of water immediately. If irritation persists.

If on skin: Wash with plenty of soap and water immediately. If irritation persists.

For medical emergencies involving this product, call for free 1-800-441-3637.

ENVIRONMENTAL HAZARDS

Do not apply this product to water, wet areas, or to any surface where water may be present. Do not apply to water.

IMPORTANT—Injury to or loss of desirable trees or other plants may result from failure to observe the following. Do not apply (except as recommended for crop use) or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, tennis courts, or similar areas. Prevent drift of dry powder or spray to desirable plants. Do not contaminate domestic waters. Keep from contact with fertilizers, insecticides, fungicides, and seeds. Thoroughly clean all traces of Hyvar X from application equipment immediately after use. Flush tank, pump, hoses, and boom with several changes of water after removing nozzle tips and screens (clean these parts separately).

Net 4 lb

© 1985 E. I. du Pont de Nemours & Co. (Inc.)

E. I. du Pont de Nemours & Co. (Inc.)
Agricultural Products Dept., Wilmington, Delaware

ACCEPTED

Aug - 3 1985

GENERAL INFORMATION

Hyvar X is a herbicide for use with water. It is a contact herbicide and is effective against a wide range of weeds and grasses. It is also effective against some types of trees and shrubs.

Hyvar X is a contact herbicide and is effective against a wide range of weeds and grasses. It is also effective against some types of trees and shrubs. It is a contact herbicide and is effective against a wide range of weeds and grasses. It is also effective against some types of trees and shrubs.

Hyvar X is a contact herbicide and is effective against a wide range of weeds and grasses. It is also effective against some types of trees and shrubs. It is a contact herbicide and is effective against a wide range of weeds and grasses. It is also effective against some types of trees and shrubs.

NOTICE OF WARRANTY

DuPont warrants that this product will perform as described on the label. It does not warrant that the product will perform as described on the label. It does not warrant that the product will perform as described on the label. It does not warrant that the product will perform as described on the label. It does not warrant that the product will perform as described on the label.

DIRECTIONS FOR USE

For application, see label. Do not apply to water, wet areas, or to any surface where water may be present. Do not apply to water.

Hyvar X is a herbicide for use with water. It is a contact herbicide and is effective against a wide range of weeds and grasses. It is also effective against some types of trees and shrubs.

DuPont will not be responsible for any damage or injury resulting from use of this product in a manner not recommended by DuPont. Do not use in a manner not recommended by DuPont.

Farm workers performing hand labor should not enter treated areas within 24 hours of application. Do not enter treated areas within 24 hours of application. Do not enter treated areas within 24 hours of application.

Written or oral warning must be given to all workers performing hand labor in or near treated areas. Do not enter treated areas within 24 hours of application. Do not enter treated areas within 24 hours of application.

Apply Hyvar X as a spray. Do not apply to water, wet areas, or to any surface where water may be present. Do not apply to water.

Before spraying, wash tank and equipment with water. Do not apply to water, wet areas, or to any surface where water may be present. Do not apply to water.

For crop use, apply with a spray. Do not apply to water, wet areas, or to any surface where water may be present. Do not apply to water.

Hyvar X is a herbicide for use with water. It is a contact herbicide and is effective against a wide range of weeds and grasses. It is also effective against some types of trees and shrubs.

[illegible]

DIRECTIONS FOR USE

Apply Hyvar X as a pre-emergent herbicide to control annual and perennial weeds in agricultural crops, lawns, and ornamental plants. Apply Hyvar X to the soil at the rate of 1 to 2 lbs. per acre. Hyvar X is a pre-emergent herbicide and must be applied before the weeds emerge from the soil. Hyvar X is not effective against weeds that are already growing in the soil.

NON-CROP USE

WEED CONTROL

Apply Hyvar X as a pre-emergent herbicide to control annual and perennial weeds in agricultural crops, lawns, and ornamental plants. Apply Hyvar X to the soil at the rate of 1 to 2 lbs. per acre.

Apply Hyvar X as a pre-emergent herbicide to control annual and perennial weeds in agricultural crops, lawns, and ornamental plants. Apply Hyvar X to the soil at the rate of 1 to 2 lbs. per acre.

Apply Hyvar X as a pre-emergent herbicide to control annual and perennial weeds in agricultural crops, lawns, and ornamental plants. Apply Hyvar X to the soil at the rate of 1 to 2 lbs. per acre.

Note: Hyvar X is not effective against weeds that are already growing in the soil.

Retreatment: Apply Hyvar X as a pre-emergent herbicide to control annual and perennial weeds in agricultural crops, lawns, and ornamental plants. Apply Hyvar X to the soil at the rate of 1 to 2 lbs. per acre.

For Small Areas: Apply Hyvar X as a pre-emergent herbicide to control annual and perennial weeds in agricultural crops, lawns, and ornamental plants. Apply Hyvar X to the soil at the rate of 1 to 2 lbs. per acre.

BRUSH CONTROL

Apply Hyvar X as a pre-emergent herbicide to control annual and perennial weeds in agricultural crops, lawns, and ornamental plants. Apply Hyvar X to the soil at the rate of 1 to 2 lbs. per acre.

Apply Hyvar X as a pre-emergent herbicide to control annual and perennial weeds in agricultural crops, lawns, and ornamental plants. Apply Hyvar X to the soil at the rate of 1 to 2 lbs. per acre.

Broadcast Treatment: Apply Hyvar X as a pre-emergent herbicide to control annual and perennial weeds in agricultural crops, lawns, and ornamental plants. Apply Hyvar X to the soil at the rate of 1 to 2 lbs. per acre.

Basal (Spot) Treatment: Apply Hyvar X as a pre-emergent herbicide to control annual and perennial weeds in agricultural crops, lawns, and ornamental plants. Apply Hyvar X to the soil at the rate of 1 to 2 lbs. per acre.

SELECTIVE USE IN CROPS

All dosages of Hyvar X are expressed as broadcast rates. For treatment use proportionately less, for example, use 1/2 of the broadcast rate when band treating 1/2 of the area.

Moisture is necessary to activate the chemical. Best results are obtained if moisture is supplied by rainfall or irrigation within two weeks after application.

CITRUS

Apply a band or broadcast treatment beneath and between trees. Avoid contact of foliage and fruit with spray or mist. Temporary yellowing of citrus leaves may occur following treatment. Do not use on citrus in Kern County, California.

Herbicide injury to citrus trees may result if citrus trees are planted in areas with high water table or poorly drained soils. Do not apply more than 8 lbs. per acre per year. Do not treat trees planted in irrigation furrows. Do not treat diseased trees such as those with citrus greening. Do not use on citrus orchards interplanted with other trees or desirable plants. Do not use on citrus plantings or orchards where citrus trees are planted in rows. Do not use on citrus trees planted in rows. Do not use on citrus trees planted in rows.

Trees Established for Four Years or More

Annual Weeds: Including barnyardgrass, watergrass, crabgrass, proso millet, Florida pusley, henbit, lambsquarters, mustard, natagrass, redtop, puncturevine, purslane, sandspur, sedge annual, sprangletop, Texas panicum, Colorado grass, and ticklebug. Apply 2 to 4 lbs. per acre. Apply once a year, preferably shortly before or after weed growth begins when adequate moisture is available.

Perennial Weeds: Best results are obtained if application is made shortly before or shortly after weed growth begins if top growth is present. Remove weeds and spray the ground. Effects on perennial weeds are slow to appear, usually progressing over a period of several months.

Make a single application per year during the period from winter to early summer. Use at the following rates:

Soil Texture	Lbs. Hyvar X Per Acre
Sand, heavy sand	4 to 5
Sandy loam	5 to 6
Silt loam, clay loam	6 to 8

Alternatively, make two applications of 3 to 4 lbs. Hyvar X per acre per year. In Florida, Texas, and Louisiana, apply in spring and summer. In California and Arizona, apply in fall and spring.

NOTICE TO BUYER

Note: For the control of weeds in citrus, the best treatment is a pre-emergent herbicide. Hyvar X is a pre-emergent herbicide and must be applied before the weeds emerge from the soil.

Perennial Weeds	Annual Weeds
Crabgrass, proso millet, Florida pusley, henbit, lambsquarters, mustard, natagrass, redtop, puncturevine, purslane, sandspur, sedge annual, sprangletop, Texas panicum, Colorado grass, and ticklebug.	Crabgrass, proso millet, Florida pusley, henbit, lambsquarters, mustard, natagrass, redtop, puncturevine, purslane, sandspur, sedge annual, sprangletop, Texas panicum, Colorado grass, and ticklebug.

Hyvar X is a pre-emergent herbicide and must be applied before the weeds emerge from the soil. Hyvar X is not effective against weeds that are already growing in the soil.

Florida Only—Trees Established Less Than Four Years

Trees Established Less Than One Year: Apply 2 to 4 lbs. per acre as needed for weed control. Do not apply at intervals less than six days. Do not apply more than 6 lbs. per acre within any 6-month period. Do not apply more than 10 lbs. per acre during the first year.

Trees Established One to Four Years: Apply 2 to 4 lbs. per acre as needed for weed control. Do not apply at intervals less than six days. Do not apply more than 6 lbs. per acre within any 6-month period. Do not apply more than 10 lbs. per acre during the first year.

Torpedograss Control — Barrier Strip Treatment

For control of torpedograss adjacent to citrus groves, apply 50 lbs. per acre. Treat a border strip 10 to 20 ft. wide adjacent to the citrus grove. Do not apply more than 100 lbs. per acre. Do not apply more than 100 lbs. per acre. Do not apply more than 100 lbs. per acre. Do not apply more than 100 lbs. per acre.

PINEAPPLE

Do not replant treated areas for any crop other than pineapple within 2 years after last application as injury to subsequent crops may result.

Hawaii and Florida: For control of seedling weeds such as amaranthus, basmatia, choris, crabgrass, Florida crabgrass, foxtail, goosegrass, (weirgrass), and Hairy apply 2 to 6 lbs. per acre broadcast before the planting material begins to grow. Use the lower rates in low rainfall areas (5 to 10 inches annually) and on clean culture fields. Use the higher rates in high rainfall areas (above 10 inches annually) and for trashmuck fields. Follow with 2 to 4 lbs. per acre as directed. Interline spray as needed prior to differentiation or apply 2 lbs. per acre broadcast after differentiation. Do not apply more than 15 lbs. total per acre per planting crop. For ratoon crop, apply 1 to 4 lbs. per acre broadcast after harvesting plant crop but before differentiation. Do not apply more than 4 lbs. total per acre per ratoon crop.

Puerto Rico: For control of seedling weeds such as crabgrass, goosegrass, (weirgrass), pigweed, and purslane, apply 2 to 4 lbs. per acre broadcast immediately after planting and before planting material begins to grow.

NOTICE TO BUYER

Purchase of this material does not confer any rights under patents of countries outside of the United States.

STORAGE AND DISPOSAL

STORAGE: Store product in original container only, away from other pesticides, fertilizer, food, or feed.

DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.