



MANZATE I



MANZATE I

WITH A ZINC SALT ADDED

ACTIVE INGREDIENT:
Manzate (Manganese ethylenebiscarbamate) 80%
(Manganese equivalent as metal) . . . 16.0%

INERT INGREDIENTS: 20%
U.S. Pat. 2,782,821-2,782,822 U.S. Pat. 2,872,333

CAUTION! KEEP AWAY FROM EYES, NOSE, THROAT, AND SKIN.
MAY BE HARMFUL IF INHALED OR SWALLOWED.
Avoid breathing dust or spray mist.
Avoid contact with skin, eyes, and clothing.
Keep away from fire or sparks.

In case of contact: Wash skin or eyes with plenty of water. For eyes, get medical attention.
WASH AWAY!—Never allow "Manzate" I to become wet during storage. This may lead to certain chemical changes which will reduce the effectiveness of "Manzate" I as a fungicide and create vapors which may be harmful. Keep container closed when not in use.

GENERAL INFORMATION:
The Fungicide "Manzate" I is recommended for use on plants for the control of various diseases (Oomycetes, rusts, and for the control of crown and collar rot) on a wide variety of plants including: *Glomerales*, *Ascomycetes*, *Basidiomycetes*, and *Zygomycetes*.
Material must be kept in suspension at all times by continuous agitation. Agitate by mechanical or hydraulic means.
Do not use container. Empty when empty. This product is toxic to fish. Do not dispose of any of this product or container in or near any body of water.

DIRECTIONS:
BRASSICA CRUCIFERAE (CABBAGES, BRUSSELS SPROUTS, ETC.): Apply as a spray at the rate of 3 to 5 lb. Manzate I per acre in sufficient water (or other suitable carrier) to thoroughly cover all plant surfaces. Begin application when symptoms indicate a disease and repeat at 7 week intervals. To prepare spray, wash required amount of "Manzate" I through the filter screen into spray tank or supply tank as being filled and use in carrying tank with "Manzate" I added after "Manzate" I is thoroughly mixed with water and when tank is half full. Allow mixture to become thoroughly emulsified. Then add the remainder of water. Continue agitation if all tank has been sprayed out.
CROWN ROT AND COLLAR ROT (CELERY, ETC.): Apply as a spray to the crown and root at the rate of 2 to 4 lbs. per 100 gals. water. Use 5 gals. of the spray mixture to treat approximately 100 sq. ft. of fruit. Apply following harvest of fruit as the fruit is passed through a high speed or clean water-rinsing spray and the fungus treatment should follow within three conditions. To prepare spray, wash required amount of "Manzate" I through the filter screen into the spray tank or supply tank in carrying tank. Continue agitation if tank has been sprayed out.

NET 50LBS.

E. I. DU PONT DE NEMOURS & COMPANY (INC.)
WILMINGTON, DELAWARE

VEA ETIQUETA EN ESPAÑOL EN EL LADO OPUESTO

THIS BAG COATED WITH
DU PONT LUDOX COLLOIDAL SILICA
TO PROVIDE ANTI-HELD CHARACTERISTICS



MANZATE I

SPRAYER LABEL

ACCEPTED
JAN 4 1969
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND HERBICIDE ACT
FOR ECONOMIC POISON REGISTER
ED UNDER NO. 352-306



MANZATE T

WITH A ZINC SALT ADDED

INGREDIENTS:
Fungicide equivalent to malathion 88%
(Fungicide equivalent to malathion 14.9%)
Inert 20%
Net Weight 50 lbs. (22.68 kg.)

PRECAUTION: Keep out of reach of children.
Avoid contact with eyes, nose, throat, and skin.
Wash hands thoroughly with soap and water after use.

GENERAL INFORMATION: Manzate T is a fungicide...
It is used to control various plant diseases...
It is effective against a wide range of fungi...

GENERAL INFORMATION: Manzate T is a fungicide...
It is used to control various plant diseases...
It is effective against a wide range of fungi...
DIRECCIONES: Manzate T es un fungicida...
Se utiliza para controlar diversas enfermedades...
Es efectivo contra una amplia gama de hongos...

NET 50LBS.

E. I. DU PONT DE NEMOURS & COMPANY (INC.)
WILMINGTON, DELAWARE, U.S.A.

SEE OPPOSITE SIDE FOR ENGLISH LANGUAGE LABEL

THIS BAG COATED WITH
DU PONT LUBOX COLLOIDAL SILICA
TO PROVIDE ANTI-SKID CHARACTERISTICS



MANZATE T

SPRAYER LABEL



MANZATE T

CON SAL DE ZINC

INGREDIENTE ACTIVO:
Manzate (Fungicida equivalente a malatión) 88.0%
(Fungicida equivalente a malatión) 14.9%
Inertes 20.0%
Peso Neto 50 libras (22.68 kg.)

PRECAUCIÓN: Manténgase fuera del alcance de los niños.
Evite el contacto con los ojos, nariz, boca y piel.
Lávese las manos con agua y jabón después de usarlo.

INFORMACION GENERAL: Manzate T es un fungicida...
Se utiliza para controlar diversas enfermedades...
Es efectivo contra una amplia gama de hongos...

INFORMACION GENERAL: Manzate T es un fungicida...
Se utiliza para controlar diversas enfermedades...
Es efectivo contra una amplia gama de hongos...
DIRECCIONES: Manzate T es un fungicida...
Se utiliza para controlar diversas enfermedades...
Es efectivo contra una amplia gama de hongos...

PESO NETO: 22,68 KILOS (50 LIBRAS)

E. I. DU PONT DE NEMOURS & COMPANY (INC.)
WILMINGTON, DELAWARE, U.S.A.

SEE OPPOSITE SIDE FOR ENGLISH LANGUAGE LABEL

PARA PROPORCIONAR RESISTENCIA AL DESGASTE
ESTA BOLSA HA SIDO TRATADA CON "LUBOX"
SILICE COLLOIDAL DE DU PONT.

DU PONT

REG. U. S. PAT. OFF.

MANZATE T

WITH A ZINC SALT ADDED

ACTIVE INGREDIENT:
 Maneb (Manganese ethylenebisdithiocarbamate)..... 80%
 (Manganese equivalent as metallic..... 16.5%)
INERT INGREDIENTS..... 20%
 U.S. Pat. 2,718,820-2,779,610 USDA Reg. No. 352-306

Keep out of reach of children.

CAUTION! MAY IRRITATE EYES, NOSE, THROAT,
 AND SKIN.
 MAY BE HARMFUL IF INHALED OR
 SWALLOWED.

Avoid breathing dust or spray mist.
 Avoid contact with skin, eyes, and clothing.
 Keep away from fire or sparks.

In case of contact, flush skin or eyes with plenty of water; for eyes, get medical attention.

IMPORTANT—Never allow "Manzate" T to become wet during storage. This may lead to certain chemical changes which will reduce the effectiveness of "Manzate" T as a fungicide and create vapors which may be flammable. Keep container closed when not in use.

GENERAL INFORMATION

Du Pont "Manzate" T Maneb Fungicide is recommended for use on bananas for the control of Sigatoka disease (*Cercospora musseana*), and for the post-harvest control of crown rot (*Phytophthora* spp.) and other diseases including *Gloeosporium musseana*, *Phytophthora perniciosa* and *Fusicladium* spp.)

Material must be kept in suspension at all times by continuous agitation. Agitate by mechanical or hydraulic means.

Do not re-use container. Bury when empty. This product is toxic to fish. Do not dispose of any of this product or container in or near any body of water.

DIRECTIONS

SIGATOKA DISEASE (CERCOSPORA MUSSEANA): Apply as a spray at the rate of 3 to 5 lbs. "Manzate" T per acre in sufficient water (or other suitable carrier) to thoroughly cover all plant surfaces. Begin applications when Sigatoka incidence is apparent, and repeat at 2 week intervals. To prepare spray, wash required amount of "Manzate" T through the filler screen while spray tank or supply tank is being filled. If oil is used in combination with "Manzate" T, add oil after "Manzate" T is thoroughly mixed with water and when tank is half full. Allow mixture to become thoroughly emulsified; then add the remainder of water. Continue agitation until tank has been sprayed out.

CROWN ROT AND SURFACE MOLDS: Apply as a spray to the crowns and fruit at the rate of 2 to 4 lbs. per 100 gals. water. Use 5 gals. of the spray mixture to treat approximately 100 trays of fruit. Apply following harvest of bananas. The fruit is passed through a hydrocooler or clean water-rinse spray, and the fungicide treatment should follow either of these procedures. To prepare spray, wash required amount of "Manzate" T through the filler screen while the spray tank or supply tank is being filled. Continue agitation until tank has been sprayed out.

NET 50 LBS.

E. I. DU PONT DE NEMOURS & COMPANY (INC.)
 INDUSTRIAL AND ENGINEERING DEPT. WILMINGTON, DELAWARE—U.S.A.

968

VEA ETIQUETA EN ESPAÑOL EN EL LADO OPUESTO

DU PONT

MANZATE T

MANZATE T

SPECIAL LABEL

SPECIAL LABEL

WAVZATE I

CON SAL DE ZINC

INGREDIENTE ACTIVO

Salado (sulfato de zinc) de manganeso) 20.0%

(Equivalentes en manganeso metálico 14.9%)

INGREDIENTES INERTES

..... 20.0%

Manufacturado fuera del alcance de las leyes.

IPRECAUCION!

¡MANTENER FUERA DEL ALCANCE DE LOS NIÑOS.
MANTENER FUERA DEL ALCANCE DE LOS NIÑOS.
MANTENER FUERA DEL ALCANCE DE LOS NIÑOS.

Evitar el contacto con la piel, ojos y ropa.

Manténgase lejos del fuego o de las chispas.

En caso de haber contacto, lavea la piel o los ojos con agua abundante para los ojos
valer a un médico.

¡IMPORTANTE!—Nunca debe permitirse que el "Wavzate I" se moje mientras está almacenado. Solo puede conducir a ciertos cambios químicos que reducirán la eficiencia del "Wavzate I" como fungicida y formar gases que pueden ser inflamables. Manténgase el contenedor cerrado cuando no se está usando.

INFORMACION GENERAL

El Fungicida General "Wavzate I" de Du Pont es recomendado en tanques para el control de la roya (patotrozo marrón) y después de la cosecha para la protección contra la roya y la pudrición de las raíces (causada por organismos tales como *Botrytis cinerea*, *Phytophthora*, *Phoma* y *Phytophthora* spp.).

Este material deberá mantenerse en suspensión en todo momento por medio de ciertos agitados mecánicos o hidráulicos.

No vuelva a usar el spray. Evitarlo cuando se vacía. Este producto es dañino a los peces. No se debe usar este producto o su envase en, o cerca de, ninguna estera de agua.

DIRECCIONES

SALENTINA (CERCA DEL MAR): Aplíquese "Wavzate I" en forma de rocio a la dosis de 3 a 5 lbs por acre en un sistema cerrado de agua (u otro sistema apropiado) para cubrir uniformemente toda la superficie de las plantas. Emplearse las aplicaciones cuando aparece la enfermedad y replantarse a intervalos de 2 semanas. Para preparar el rocio, hágase pasar la cantidad necesaria del producto a través del tanque mientras se llena el tanque. Si se usa agua en combinación con "Wavzate I", añádase el aceite después que el "Wavzate I" está bien mezclado con el agua y cuando el tanque se encuentra lleno hasta la mitad. Permítase que la mezcla se estabilice bien; entonces añádase el resto del agua. Dado que el agua se agita hasta que el tanque se encuentra en 3/4 de su capacidad, asegúrese de que el tanque se encuentre en 3/4 de su capacidad.

PODCALCANTO (CERCA DEL MAR): Aplíquese "Wavzate I" en forma de rocio a las coronas y follaje a la dosis de 2 a 4 lbs por 100 gal. de agua. Use entre 5 y 7 gal. de rocio para tratar aproximadamente 100 coronas de follaje. Aplíquese después de la cosecha de los bananos. Si para la fruta a través de un hidro-sprayer o cualquier tipo de rocio, y el tratamiento tardado deberá hacerse inmediatamente después de uno de estos procedimientos. Para preparar el rocio, hágase pasar la cantidad necesaria del producto a través del tanque mientras se llena el tanque. Dado que el agua se agita hasta que el tanque se encuentra en 3/4 de su capacidad, asegúrese de que el tanque se encuentre en 3/4 de su capacidad.

PESO NETO: 22,68 KILOS (50 LIBRAS)

E. I. DU PONT DE NEMOURS & COMPANY (INC.)
INDUSTRIAL AND BIOTECHNICALS DEPT., WILMINGTON, DELAWARE, U.S.A.

SEE OPPOSITE SIDE FOR ENGLISH LANGUAGE LABEL