

DIRECTIONS

Cotton – Beans – Soybeans – Sugar Beets

SUPPLEMENTAL SEED TREATMENT—Apply "Demosan" 65W as a slurry at the rate of 6 ozs. (West of Rocky Mountains, use 10 ozs.) per 100 lbs. of acid delinted, reginned. or fuzzy cottonseed; 4 ozs. per 100 lbs. of beans and soybeans; and 6 ozs. per 100 Ibs. of sugar beets. "Demosan" 65W may be applied separately as a seed overcoat or in combination with Du Pont seed protectants such as "Arasan" 70-S or "Arasan" 42-S.

SEED OVERCOATING: For standard slurry treaters mix "Demosan" 65W in water and apply at the rates, slurry cup sizes and dump

ghts shown below. For the "Panogen"* and "Mist-O-Matic"* treaters equipped to apply slurries, adjust slurry cups and dump weights to apply the amount of slurry (cc per 100 lbs. of seed) shown. One fluid ounce is equivalent to 29.5 cc.

| Crop Seed | Treating Rate Oz./Cwt. | Type of Slurry Treater | Concentra- tion Lbs./ Gal. Water | Slurry cc/Cwt. Seed | Slurry Cup cc | Lbs. Dump Weight |
|----------------------|------------------------------|------------------------------|--|---------------------------|------------------|------------------------|
| Cotton | 6 | Standard | 4 | 460 | 46 | 10 |
| (East of Rockies) | | Other | 4 | 460 | | |
| Cotton | 10 | Standard | 4 | 766 | 46 | 6 |
| (West of Rockies) | | Other | 4 | 766 | | |
| Beans and | 4 | Standard | 4 | 306 | 23 | 7.5 |
| Soybeans | | Other | 4 | 306 | | |
| Sugar Beets | 6 | Standard | 4 | 460 | 46 | 10 |
| | | Other | 4 | 460 | | |

COMBINATION TREATMENT: "Demosan" 65W may be mixed in the slurry tank with "Arasan" 70-S or "Arasan" 42-S seed protectants by adjusting slurry cup sizes and dump weights for the

"Panogen" trademark of Nor-Am Agricultural Products. Inc.

"Mist-O-Matic" trademark of Gustafson Manufacturing Co.

CONTINUED ON RIGHT PANEL



ACTIVE INGREDIENT: Chloroneb (1,4-dichloro-2,5-dimethoxybenzene)......65%

U.S. Pat. 3,265,564 EPA Est. 97/9-MS-1

Keep out of reach of children.

CAUTION ! MAY IRRITATE EYES, NOSE, THROAT, AND SKIN.


Avoid breathing dust or spray mist. Avoid contact with skin, eyes, and clothing.



B-20709 8-74

E. I. DUPONT DENEMOURS & COMPANY (INC.) BIOCHEMICALS DEPARTMENT, WILMINGTON, DELAWARE





EPA Reg. No. 352-312-AA

When used as directed, "Demosan" 65W can be expected to inprove stands and plant vigor through the suppression of seedling olights, soreshin, and pre- and post-emergence damp-off caused by su h organisms as Rhizoctonia and Pythium species and Sclerotium rol ii. It is highly effective in preventing post-emergence damp-off, e ecially when used as a supplemental seed coating.

GENERAL INFORMATION --- Du Pont "Demosan" 65W Chioroneb

Fungicide is recommended for use as a supplemental seed treatment,

or as an in-furrow soil treatment at planting time, for the control of ceedling diseases of cotton, beans and soybeans; and as a supple-

mental seed treatment for sugar beets. It is to be used in conjunction with a standard fungicide seed treatment, such as [] Pont

"Arasan" 70-S or "Arasan" 42-S seed protectants.

MADE IN U.S.A.

PRINTED IF U.S.A.



MATION — Du Pont "Demosan" 65W Chloroneb nmended for use as a supplemental seed treatment, i soil treatment at planting time, for the control of of cotton, beans and soybeans; and as a supplement for sugar beets. It is to be used in conjuncdard fungicide seed treatment, such as Du Pont "Arasan" 42-S seed protectants.

ected, "Demosan" 65W can be expected to improve vigor through the suppression of seedling blights, - and post-emergence damp-off caused by such orgaonia and Pythium species and Sclerotium rolfsii. It is n preventing post-emergence damp-off, especially upplemental seed coating.

PRINTED IN U.S.A.



DIRECTIONS CONTINUED FROM LEFT PANEL

increased amount of water needed to handle the additional solids. Observe all cautions and limitations on labeling of all products used in mixtures.

Label Treated Seed: "This seed treated with chloroneb fungicide ('Demosan' 65W). Do not use for food, feed, or oil purposes." Add appropriate information for any other chemical treatments.

IN-FURROW TREATMENT—Apply "Demosan" 65W at the following rates as an in-furrow spray at planting time, using 10 to 20 gals. water per acre.

| | Lbs. "Demosan" 65W Per Acre (12,000 to 14,000 Row Feet) |
|--------------------|--|
| Cotton | 2 to 3 |
| Beans and Soybears | 11/2 |

Use a pressure sprayer equipped with dual row nozzles, one nozzle directed downward to the seed in the open furrow and the other backward into the covering soil. Nozzle orifice size should not be smaller than 8002 T-jet or equivalent.

NOTE: Do not use treated seed for food, feed, or oil purposes. Do not graze treated beans or soybeans within 45 days of planting. Do not re-use container; crush and bury when empty.

NOTICE TO BUYER—Seller warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. This warranty does not extend to use of this product contrary to label use directions, or under abnormal use conditions, or under conditions not reasonably foreseeable to seller; buyer assumes all risk of any such use. Seller makes no other warranties, express or implied.

Purchase of this material does not confer any rights under patents of countries outside of the United States.