




AGRICULTURAL BULLETIN



SINBAR® WEED KILLER
STRAWBERRIES



SUPPLEMENTAL LABELING
EPA REG. NO. 352-317

SINBAR® WEED KILLER FOR WEED CONTROL IN STRAWBERRIES

Du Pont Sinbar® Weed Killer is recommended for pre- and postemergence weed control in established strawberries grown in the matted row system. Sinbar® controls crabgrass (large, smooth), chickweed (common, mouse-ear), horseweed (mare's tail), lambsquarters (common), and pigweed (prostrate, redroot) and gives partial control of orchardgrass and quackgrass.

HOW TO USE

U.S. except California and Florida: Use only on plants established in the field for at least 6 months (matted row system). Make a single broadcast application of 1/2 to 1 1/4 lbs. per acre; use the lower rate on coarse-textured soils. Apply immediately after postharvest renovation (complete removal of old leaves is essential) but before new growth begins; or apply in the dormant season from fall to late winter. Treat before weeds are 2" tall or across.

Apply with a tractor-mounted, fixed-boom sprayer properly calibrated to a constant speed and rate of delivery.

Treatment may reduce runner production or plant stand. Varieties such as Guardian, Darrow, and Honeoye may be sensitive to Sinbar®; determine varietal tolerance prior to adoption as a field practice to prevent possible crop injury. Do not use on soils less than 2% organic matter and do not apply over top of newly planted strawberries as crop injury may result. Alfalfa, apples, blueberries, citrus, mint, peaches, strawberries, and sugarcane may be planted 1 year after last application. Do not plant any other crops within 2 years after application as injury to the crop may occur.

IMPORTANT

BEFORE USING SINBAR®, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on the package label. Follow the instructions carefully.

6/23/82

JUN 24 1982