



Reg # 352-317

PM-25

121

SUPPLEMENTAL LABELING

AGRICULTURAL PRODUCTS

..... A Growing Partnership With Nature



SINBAR® HERBICIDE  
WEED CONTROL IN  
STRAWBERRIES

**SINBAR® HERBICIDE**

EPA REG. NO. 352-317

**FOR WEED CONTROL IN STRAWBERRIES  
(EXCEPT CALIFORNIA AND FLORIDA)**

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with the labeling.

**HOW TO USE**

Use only on plants established in the field for at least 6 months (matted-row system)

Apply 2 - 6 ozs. product per acre per application. Make an initial application no earlier than 6 months after planting, or after postharvest renovation before new growth begins in midsummer. Applications of 2 - 6 ozs. may be made in late fall for additional control of winter annual weeds. For best results, weeds should be less than 1 inch in height or diameter when treated.

Do not apply more than 8 ozs. of "Sinbar" 80WP per acre per growing season.

**Notes:**

Do not use on soils with less than 2% organic matter.

Do not apply over the top of newly planted strawberries, as crop injury may result.

Treatment may reduce runner production or plant stand.

Avoid overlapping with spray equipment, as crop injury may result.

Do not use in the states of California and Florida.

A reduction in residual weed control may occur under adverse environmental conditions such as heavy rainfall or when low rates are used on soil high in organic matter.

**VARIETIES**

Certain varieties differ in their sensitivity to "Sinbar." Prior to adoption as a field practice, determine the varietal tolerance under expected field conditions to avoid the potential for widespread injury.

**REPLANTING (FOLLOWING STRAWBERRIES)**


Alfalfa, apples, blueberries, mint, peaches, and strawberries may be planted one year after last application. Do not plant any other crop within two years after application, as injury to the crop may occur.

**IMPORTANT - BEFORE USING "SINBAR," READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL**

This bulletin contains new or supplemental instructions for use of this product which may not appear on the package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

083091/011592/013192





PM 25

352-317

SUPPLEMENTAL LABELING

1 of 1

AGRICULTURAL PRODUCTS

'..... A Growing Partnership With Nature'



SINBAR® HERBICIDE  
WEED CONTROL IN  
STRAWBERRIES

SINBAR® HERBICIDE

EPA REG. NO. 352-317

FOR WEED CONTROL IN STRAWBERRIES  
(EXCEPT CALIFORNIA AND FLORIDA)

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with this labeling.

HOW TO USE

Use only on plants established in the field for at least 6 months (matted-row system)

Apply 2 - 6 ozs. product per acre per application. Make an initial application no earlier than 6 months after planting, or after postharvest renovation before new growth begins in midsummer. Applications of 2 - 6 ozs. may be made in late fall for additional control of winter annual weeds. For best results, weeds should be less than 1 inch in height or diameter when treated.

Do not apply more than 8 ozs. of "Sinbar" 80WP per acre per growing season.

Notes:

Do not use on soils with less than 2% organic matter.

Do not apply over the top of newly planted strawberries, as crop injury may result.

Treatment may reduce runner production or plant stand.

Avoid overlapping with spray equipment, as crop injury may result.

Do not use in the states of California and Florida.

A reduction in residual weed control may occur under adverse environmental conditions such as heavy rainfall or when low rates are used on soil high in organic matter.

VARIETIES

Certain varieties differ in their sensitivity to "Sinbar." Prior to adoption as a field practice, determine the varietal tolerance under expected field conditions to avoid the potential for widespread injury.

REPLANTING (FOLLOWING STRAWBERRIES)

Alfalfa, apples, blueberries, mint, peaches, and strawberries may be planted one year after last application. Do not plant any other crop within two years after application, as injury to the crop may occur.

IMPORTANT - BEFORE USING "SINBAR," READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on the package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

083091/011592/013192

**BEST AVAILABLE COPY**

