



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

AUG - 7 2006

Ms. Anna M. Stoops
E.I. DuPont de Nemours & Company
DuPont Crop Protection
Stine-Haskell Research Center
P.O. Box 30
Newark, DE 19714-0030

Dear Ms. Stoops:

Subject: DuPont Sinbar Herbicide (Add Watermelons)
EPA Registration No. 352-317
Application Dated July 14, 2006

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended is acceptable. A stamped copy of labeling is enclosed for your records.

Supplemental labeling must be incorporated into a master label and copies of master labeling submitted to the Agency for our files at your next printing or within two years from the date of acceptance of supplemental labeling, whichever comes first. The Agency will consider a convincing argument as to why a supplemental label should not appear on a master label on a case by case basis.

Sincerely,

James A. Tompkins
James A. Tompkins
Product Manager 25
Herbicide Branch
Registration Division (7505P)

272

SUPPLEMENTAL LABELING

**DuPont Crop
Protection**

**DUPONT™ SINBAR® HERBICIDE
FOR WEED CONTROL
IN WATERMELONS**

DUPONT™ SINBAR® HERBICIDE

EPA Reg. No. 352-317

**FOR USE OF SINBAR® ON WATERMELON
TO CONTROL BROADLEAF WEEDS**

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Dupont™ SINBAR® herbicide should be used for control of broadleaf weeds (i.e. annual morningglory) in watermelons.

A maximum of one preemergence ground application of SINBAR® (active ingredient - terbacil) may be applied at a rate of 2 to 4 ounces product per acre (0.1 to 0.2 pound active ingredient per acre). Do not exceed 4 ounces of product per acre per year (0.2 pound active ingredient per acre per year). A 70-day preharvest interval (PHI) is required.

SINBAR® may be used for direct seeded or transplant watermelons. Where seeded watermelons are used as cultural practice, make applications after planting, but before the crop emerges. Where transplants are used as a cultural practice, apply SINBAR® pre-transplant.

Where plastic mulch is used as a cultural practice, SINBAR® may be applied pre emergence under plastic mulch or to row middles.

Use the lower rate on coarse textured soils lower in organic matter.

PRECAUTIONS

Do not apply over the top or allow spray to contact crop, as unexpected crop injury may result.

Do not apply within 70 days of harvest.

A reduction in residual weed control may occur under adverse environmental conditions such as heavy rainfall or when low rates are used on soil high in organic matter.

IMPORTANT

BEFORE USING SINBAR®, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.

This bulletin contains new or supplemental instructions for use of this product which does not appear on the package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

ACCEPTED
~~with COMMENTS~~
~~January 10, 2006~~
AUG - 7 2006
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.
352-317

DR-330 070903-2