


352-317

03/21/2007
draft label

111

**DuPont Crop
Protection**



SUPPLEMENTAL LABELING

**DUPONT™ SINBAR®
HERBICIDE**

**FOR USE ON MINT IN THE NORTHERN
CALIFORNIA COUNTIES OF LASSEN,
MODOC, SHASTA AND SISKIYOU**

DUPONT™ SINBAR® HERBICIDE

EPA Reg. No. 352-317

**FOR USE ON MINT (PEPPERMINT, SPEARMINT) IN THE
NORTHERN CALIFORNIA COUNTIES OF LASSEN, MODOC,
SHASTA AND SISKIYOU**

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

APPLICATION INFORMATION

Preemergence Application: Make a single broadcast application of 1 to 2 pounds per acre before mint emerges.

Weeds controlled in mint: Weeds controlled by SINBAR® at 1 to 2 pounds per acre are dogfennel, nightshade, henbit, chickweed, tansymustard, annual pepperweed, shepherdspurse, Jim Hill mustard and filaree. Weeds partially controlled or suppressed by SINBAR® are Russian thistle, nutsedge, annual bluegrass, barnyardgrass and false flax.

Postemergence Application: Apply 1 to 1-1/2 pounds per acre before weeds are 2 inches tall (or across) and grasses are 1 inch tall (or across). Add 1/2 to 1 pint Surfactant to each 25 gallons of spray: non-phytotoxic superior-type spray oil may be substituted at the rate of 1 gallon per acre. If preemergence treatment has not been applied, a second postemergence application may be made but do not exceed 2 pounds total per acre per season. Do not apply within 60 days of harvest.

Preemergence + Postemergence: Apply as directed above. Do not apply more than 2 pounds total per acre per season. Do not apply within 60 days of harvest.

Note: For either newly planted roots or established mint, soil must be well prepared before preemergence application. Crop injury and/or poor weed control may result if application is made to ground which is cloddy or compacted, resulting in exposed or improperly covered roots. Do not apply to newly planted roots that are diseased or lacking in vigor, nor on thinly covered or exposed subsoil areas as injury to the crop may result. Use of insecticides in fields where SINBAR® is applied may result in injury to the mint; observe use limitations on insecticide labels. Treated areas may be planted to mint one year after last application.

SOIL LIMITATIONS

Crop injury may result from application of SINBAR® to sand, loamy sand or gravelly soil types or soils low in organic matter (less than 1%). For mint grown under these soil conditions, limit initial use of SINBAR® to a small area to determine crop response prior to treating larger acreage.

Checking Soil Classification and Organic Matter: Before using SINBAR, determine the soil classification and organic matter of the area of intended use. To obtain a representative soil classification and organic matter for the test area take several 0-6 inch samples from different areas of the field and analyze them separately. Consult local extension publications for additional information on recommended soil sampling procedures.

IMPORTANT

BEFORE USING SINBAR®, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

DR-677 012407-3