

DU PONT Weed Killer



LONDAX

WEED KILLER

GRANULES

ACTIVE INGREDIENTS	15%
3-(3,4-dichlorophenyl)-1-methylurea	5%
N-isopropylacetamide	10%
INERT INGREDIENTS	85%

GENERAL INFORMATION

50 lbs.

Du Pont de Nemours & Co. (Inc.)

50
LBS. NET

LONDAX G

WEED KILLER



DIRECTIONS

BROADCAST TREATMENT

BAND TREATMENT

REPLANTING

NOTE

NOTICE TO BUYER

THIS PRODUCT CONTAINS
DU PONT LUDOX COLLOIDAL SILICA
REGISTERED TRADEMARK OF DU PONT DE NEMOURS & CO.

BEST DOCUMENT AVAILABLE



LONDAX G

50 LBS. NET



LONDAX G WEED KILLER

GRANULES

ACTIVE INGREDIENTS: 15% 1,1-Dimethyl-3-methylurea 5% 2-Chloro-N-isopropylacetamide 10%

INERT INGREDIENTS: 85%

WARNING: Harmful if swallowed. May be fatal by skin absorption. Irritating to skin, eyes, nose, and throat. May cause allergic skin reaction. Do not get in eyes, on skin, or on clothing. Avoid breathing dust.

While handling, wear rubber gloves. Wash thoroughly after handling. Launder clothing before re-use. In case of contact, flush skin or eyes with plenty of water. If eyes get medical attention.

Net 50 lbs.

E. I. du Pont de Nemours & Co. (Inc.) Industrial and Biochemicals Dept., Wilmington, Delaware

IMPORTANT: Injury to or loss of desirable trees or other plants may result from failure to observe the following: Do not apply (except as recommended for crop use) or drain or drain into ground on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawn, walk, driveway, tennis courts, or similar areas. Prevent drift to desirable plants. Do not contaminate any body of water. Keep from contact with herbicides, insecticides, fungicides, and seeds. Thoroughly clean all Londax G Weed Killer application equipment immediately after use. Do not apply to a wet surface.

GENERAL INFORMATION: Du Pont "Londax" G Weed Killer is a granular product to be applied as a pre-emergence treatment for the selective control of annual weeds in soybeans (seed crop only) and field corn. It is non-corrosive to equipment, non-toxic, non-flammable. Du Pont "Londax" G Weed Killer selectively controls seedling annual weeds such as bar, pigweed, crabgrass, foxtail (excluding giant foxtail), lambsquarters, gooseweed, ragweed and barnyard. It will not control perennial weeds such as Bermudagrass, Johnsongrass, nutgrass, or quackgrass.

50 LBS. NET

LONDAX G



BEST DOCUMENT AVAILABLE

SPECIMEN LABEL

44

ACCEPTED JAN 2 1952 352-334

8% OF ACTUAL SIZE

352-334

2A

DIRECTIONS

Du Pont "Londax" G Weed Killer should be used only in accordance with recommendations on this label, or in separate published Du Pont recommendations available through local dealers.

SOYBEANS (SEED CROP ONLY) AND FIELD CORN PRE-EMERGENCE APPLICATION

Make a single application as a band or broadcast treatment after planting but before crop emerges. Apply with tractor-mounted application equipment properly calibrated to a constant speed and rate of delivery. Use at the following rates.

BROADCAST TREATMENT

Soil Type†	Lbs. "Londax" G Per Acre	
	Low organic matter (1 to 3%)	Moderate organic matter (3 to 6%)
Sandy loam	6½ to 13½	13½ to 20
Silt loam	10 to 16½	16½ to 25
Clay loam	13½ to 20	20 to 30

†Do not use on sand or loamy sand nor on soils with less than 1% organic matter as crop injury may result. Use the lower rates on soils with minimum organic matter listed and the higher rates on maximum organic matter listed.

BAND TREATMENT—Use proportionately less; for example, for 14" band on 42" row, use 1/3 of the broadcast rate.

Soil should be well prepared and as free as possible from trash and clods. Plant seed at least 1 1/4" deep on flat or raised seedbeds only or injury to the crop may result. Sufficient moisture (usually 1" to 2") in the form of rainfall or irrigation is necessary after treatment to carry the chemical into the root zone of germinating weeds; best results are obtained when this occurs within two weeks after application. If moisture is insufficient to activate the chemical or if the soil becomes crusted before crop emerges, a shallow rotary hoeing should be made before weeds become well established; otherwise, avoid cultivating or disturbing the treated area. Do not cover treated band with soil while cultivating middles.

REPLANTING—If initial seeding fails to produce a stand, soybeans (seed crop only) and field corn may be replanted in soil treated with "Londax" G Weed Killer. Thoroughly rework soil before replanting. Do not re-treat field with a second application during the same crop year as injury to the crop may result. Do not replant to other crops within 4 months after application as injury to subsequent crops may result.

NOTE—Do not graze or feed soybean forage or silage from treated areas to livestock. Do not use soybeans for food, feed or oil purposes.

NOTICE TO BUYER: Seller warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. This warranty does not extend to use of this product contrary to label use directions, or under abnormal use conditions, or under conditions not reasonably foreseeable to seller; buyer assumes all risk of any such use. Seller makes no other warranties, express or implied.

THIS BAG COATED WITH
DU PONT LUDOX® COLLOIDAL SILICA
TO PROVIDE ANTI-SEED CHARACTERISTICS

45