


(A)  **United States Environmental Protection Agency**
Office of Pesticide Programs (H7505C)
 Washington, DC 20460

Application for Pesticide:

Registration
 Amendment
 Other

OPP Identifier Number
256995

Section I

1. Company/Product Number
 352-341

2. EPA Product Manager
 Mary L. Waller

3. Proposed Classification
 None Restricted

4. Company/Product (Name)
 DuPont Manzate® 200 Fungicide

PM#
 21

5. Name and Address of Applicant (Include ZIP Code)
 E. I. du Pont de Nemours & Company
 Barley Mill Plaza, P. O. Box 80038
 Wilmington, DE 19880-0038
 Attn: Mary Lou Hawkins, WM6-126
 Check if this is a new address

6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to:
 EPA Reg. No. _____
 Product Name _____

Section II

Amendment - Explain below

Resubmission in response to Agency letter dated _____

Notification - Explain below.

Final printed labels in response to Agency letter dated _____ in accordance with PR Notice 95-2

"Me Too" Application. **APR 30 1997**

Other - explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Change via notification: add pest (gray leaf spot) to corn. New final label, SL-288-1 9116 11/18/96, is attached. This notification is consistent with the provisions of PR Notice 95-2 and EPA regulations at 40 CFR 152.45, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 95-2 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section III

1. Material This Product Will Be Packaged In:

Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted.		If "Yes," Unit Package wgt. No. per container	If "Yes," Package wgt. No. per container

3. Location of Net Contents Information
 Label Container

4. Size(s) of Retail Container

5. Location of Label Directions
 On Label
 On Labeling accompanying product

6. Manner In Which Label Is Affixed To Product
 Lithograph
 Paper glued
 Stenciled
 Other (_____)


Section IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)

Name Mary Lou Hawkins	Title Product Registration Manager	Telephone No. (Include Area Code) (302) 892-8805
--------------------------	---------------------------------------	---

Certification

I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

2. Signature


3. Title
 Product Registration Manager

4. Typed Name
 Mary Lou Hawkins

5. Date
 4/17/97

6. Date Application Received (Stamped)



MANZATE® 200

2/11

FUNGICIDE
WETTABLE POWDER
BY WEIGHT

ACTIVE INGREDIENT

A coordination product of

zinc ion and manganese ethylene bisdithiocarbamate in which the ingredients are:	80%
Manganese++	16%
Zinc++	2%
Ethylene bisdithiocarbamate ion (C ₄ H ₆ N ₂ S ₄)=	62%

INERT INGREDIENTS 20%

This product contains the toxic inert ingredient paraformaldehyde. TOTAL 100%

EPA Reg No. 352-341

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION! MAY IRRITATE EYES, NOSE, THROAT AND SKIN. MAY BE HARMFUL IF ABSORBED THROUGH SKIN. INHALED OR SWALLOWED. Avoid breathing dust or spray mist. Avoid contact with skin, eyes and clothing. Keep away from fire or sparks.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Dilute by giving two glasses of water to drink and call a physician. Never give anything by mouth to an unconscious person.

IF INHALED: Move subject to fresh air. Consult a physician.

IF IN EYES: Flush eyes with large amounts of water for at least 15 minutes. Consult a physician if irritation persists.

IF ON SKIN: Wash affected skin areas with soap and water. Consult a physician if irritation persists.

For medical emergencies involving this product, call toll free 1-800-441-3637.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers (other than mixers and loaders) must wear:

- Coveralls over long-sleeved shirt and long pants.
- Waterproof gloves.
- Shoes plus socks.

Mixers and Loaders must wear:

- Coveralls over long-sleeved shirt and long pants.
- Waterproof gloves.
- Shoes plus socks.
- Protective eyewear.
- Chemical-resistant apron when mixing or loading.

Discard clothing or other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

During aerial application, human flaggers must be in enclosed cabs.

ENGINEERING CONTROL STATEMENTS

When handlers use enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR part 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove personal protective equipment immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Cover or incorporate spilled treated seed. Do not contaminate water by disposing of equipment washwaters.

NOTIFICATION
APR 30 1997

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over long-sleeved shirt and long pants.
- Waterproof gloves.
- Shoes plus socks.

"Manzate" 200 should be used only in accordance with recommendations on this folder or in separate published Du Pont recommendations available through local dealers.

Du Pont will not be responsible for losses or damages resulting from use of this product in any manner not specifically recommended by Du Pont. User assumes all risks associated with such nonrecommended use.

"Manzate" 200, a wettable powder containing mancozeb, is recommended for use as a spray for the control of many important plant diseases.

APPLICATION INSTRUCTIONS

AS A SPRAY (Ground or Aerial Equipment)—Apply "Manzate" 200 at the rate shown; use sufficient water to provide thorough coverage, use 20 to 100 gals per acre for ground equipment and no less than 2 gals per acre for aircraft. Add "Manzate" 200 slowly to water in the spray tank with agitation, or premix thoroughly in separate holding tank for concentrate or aircraft sprayers. Continuous agitation is required to keep the product in suspension. Add a spreader-sticker spray adjuvant if needed.

RESTRICTIONS

Foliar Applications

Where EBDC Products Used Allow the Same Maximum Poundage of Active Ingredient Per Acre Per Season

If more than one product containing an EBDC active ingredient (maneb, mancozeb or metiram) is used on a crop during the same

growing season and the EBDC products used allow the same maximum poundage of active ingredient per acre per season, then the total poundage of all such EBDC products used must not exceed any one of the specified individual EBDC product maximum seasonal poundage of active ingredient allowed per acre.

Where EBDC Products Used Allow Different Maximum Poundage of Active Ingredient Per Acre Per Season

If more than one product containing an EBDC active ingredient is used on a crop during the same growing season and the EBDC products used allow different maximum poundage of active ingredient per acre per season, then the total poundage of all such EBDC products used must not exceed the lowest specified individual EBDC product maximum seasonal poundage of active ingredient allowed per acre.

Seed Treatment

In addition to the maximum number of foliar applications permitted by the formula stated above, a single application for seed treatment may be made on crops which have registered seed treatment uses.

CHEMIGATION

Apply "Manzate" 200 Fungicide only through sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set or hand move irrigation systems only on crops specifically designated in the "Directions For Use" for that particular crop. Do not apply "Manzate" 200 through any other type of irrigation system or on any other crops.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service Specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Specific Instructions for Public Water Systems:

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.
8. Good agitation is required in the injection tank.
9. In moving systems, apply specified dosage of "Manzate" 200 as a continuous injection. In nonmoving systems inject "Manzate" 200 for 15 to 30 minutes at end of cycle. Use the least amount of water possible consistent with uniform coverage.
10. Mix the amount of "Manzate" 200 needed for acreage to be treated into the quantity of water determined during prior calibration. For moving systems inject into the system continuously for one complete revolution of the field. For nonmoving systems inject into system for the time established during calibration.
11. Stop injection equipment after treatment is completed and continue to operate irrigation equipment until all "Manzate" 200 is flushed from system.

Specific Instructions for Sprinkler Irrigation Systems:

1. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the

- system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.
8. Good agitation is required in the injection tank.
9. In moving systems, apply specified dosage of "Manzate" 200 as a continuous injection. In nonmoving systems inject "Manzate" 200 for 15 to 30 minutes at end of cycle. Use the least amount of water possible consistent with uniform coverage.
10. Mix the amount of "Manzate" 200 needed for acreage to be treated into the quantity of water determined during prior calibration. For moving systems inject into the system continuously for one complete revolution of the field. For nonmoving systems inject into system for the time established during calibration.
11. Stop injection equipment after treatment is completed and continue to operate irrigation equipment until all "Manzate" 200 is flushed from system.

CROP	DISEASES CONTROLLED	RATE OF MANZATE 200 PER APPLICATION LBS/ACRE	TIMING/INTERVALS (Also refer to Directions for Use)	RESTRICTIONS/COMMENTS
Apples (See Pomefruit)				
Asparagus	Cercospora leaf spot Rust	2	Start applications when rust first appears and repeat at 10 day intervals. Four applications are usually sufficient.	Apply only on asparagus ferns after spears have been harvested. Do not apply within 180 days of harvest in all states except CA and AZ (120 days). Do not apply more than 8 lbs (6.4 lbs active) per acre per season.
Asparagus Crown (planting stock)	Crown rot	1 lb per 100 gals	Dip clean, loosely packed crowns into continuously agitated fungicide suspension for 5 minutes. Drain and plant as soon as possible.	Wash dirty crowns before dip treatment. Replace suspension in clean tank when discolored by soil.
Bananas	Sigatoka	2 to 3	Apply when leaves first appear and repeat every 14 to 21 days or as required. Use sufficient water to provide adequate coverage. The addition of a surfactant to spray solutions will improve performance.	Do not apply more than 30 lbs (24 lbs active) per acre per growing cycle. Minimum preharvest interval (0 days).

CROP	DISEASES CONTROLLED	RATE OF MANZATE 200 PER APPLICATION LBS/ACRE	TIMING/INTERVALS (Also refer to Directions for Use)	RESTRICTIONS/COMMENTS
Barley, Oats, Rye, Wheat	Helminthosporium leaf spot Leaf rust Septoria glume blotch Septoria leaf spot Tan spot	2	Start application at onset of disease or when plants are in the tillering to jointing stage and repeat at 7 to 10 day intervals. The addition of a surfactant to spray solutions will improve performance.	Do not make more than three applications during the season. Do not apply more than 6 lbs (4.8 lbs active) per acre per crop. Do not apply after Feekes growth stage 10.5 or heading, nor within 26 days of harvest. Do not graze livestock in treated areas prior to harvest.
Caprifig (Non-food use)	Endosepsis (Fusarium) Mold	4 lbs/100 gals	Prepare mamme figs by making a shallow cut through the eye and then hand dividing to avoid wasp injury. Submerge mamme figs in the continuously agitated suspension for at least 15 minutes. Drain before placement in trees.	Use fresh dipping suspension after treating 4 to 5 batches of figs.
Corn (sweet corn, popcorn and sweet corn for hybrid seed production), (field and field corn for hybrid seed production)	Common rust Helminthosporium leaf blight Gray leaf spot	1 1/2	Use sufficient water for thorough coverage. Start applications when disease first appears and repeat at 4 to 7 day intervals. The addition of a surfactant to spray solutions will improve performance.	Do not apply within 7 days of harvest. Do not apply more than 22.5 lbs (18 lbs active) per acre per crop east of Mississippi and AR and LA. Do not apply more than 7.5 lbs (6 lbs active) per acre per crop west of the Mississippi except AR and LA. Do not feed treated forage to livestock. Do not apply within 40 days of harvest. Do not apply more than 15 lbs (12 lbs active) per acre per crop. Do not feed treated forage to livestock.
Cotton (Southwest only)	Rust	1 1/4 to 2	Begin when rust first appears in the area. Repeat at 10 to 14 day intervals.	Do not apply after bolls open. Do not apply more than 8 lbs (6.4 lbs active) per acre per crop. Do not apply within 45 days of harvest. Do not feed treated forage to livestock. Do not graze livestock in treated areas. Do not feed gin trash to livestock.

CROP	DISEASES CONTROLLED	RATE OF MANZATE 200 PER APPLICATION LBS/ACRE	TIMING/INTERVALS (Also refer to Directions for Use)	RESTRICTIONS/COMMENTS
Cranberries	Fruit rot	3 to 6	Start applications at mid-bloom and repeat at 7 to 10 day intervals.	Do not apply within 30 days of harvest. Do not apply more than 18 lbs (14.4 lbs active) per acre per season.
Cucumbers	Anthracnose Cercospora leaf spot Downy mildew Gummy stem blight Scab	2 to 3	Start applications when plants are in the two-leaf stage and repeat at 7 to 10 day intervals. Use sufficient water and direct spray to provide thorough coverage of both upper and lower leaf surfaces.	Do not apply within 5 days of harvest. Do not apply more than 24 lbs (19.2 lbs active) per acre per crop.
Fennel	Early & late blights	2	Begin in plant beds at emergence. Repeat at 7 to 10 day intervals.	Do not apply within 14 days of harvest. Do not apply more than 16 lbs (12.8 lbs active) per acre per crop. Do not graze treated area to livestock.
Grapes (East of the Rocky Mountains)	Black rot Bunch rot Deadarm Downy mildew	1 1/2 to 4	Apply in sufficient water to provide thorough coverage starting when new shoots are 1/2 to 1 1/2 inches long. Repeat when shoots are 3 to 5 inches long, when shoots are 8 to 10 inches long, and then at 7 to 10 day intervals until fruit is set. For late season control of black rot, deadarm and downy mildew the use of other approved and recommended fungicides is suggested.	Do not apply within 66 days of harvest. Do not apply more than 24 lbs (19.2 lbs active) per acre per season.
(West of the Rocky Mountains)		1 1/2 to 2 1/2		Do not apply within 66 days of harvest except in CA where no application can be made after bloom. Do not apply more than 7.5 lbs (6 lbs active) per acre per season.

CROP	DISEASES CONTROLLED	RATE OF MANZATE 200 PER APPLICATION LBS/ACRE	TIMING/INTERVALS (Also refer to Directions for Use)	RESTRICTIONS/COMMENTS
Melons Cantaloupes Casaba Crenshaw Honeydew Watermelon	Alternaria leaf spot Anthracnose Downy mildew Gummy stem blight Cercospora leaf spot	2 to 3	Start applications when plants are in the two-leaf stage and repeat at 7 to 10 day intervals. Use sufficient water and direct spray to provide thorough coverage of both upper and lower leaf surfaces. Some varieties are sensitive to Manzate 200 fungicide. Consult State Cooperative Extension Service Specialist prior to use.	Do not apply within 5 days of harvest. Do not apply more than 24 lbs (19.2 lbs active) per acre per crop.
Oats (See barley)				
Onions (dry bulb) ((fallow drench))	Botrytis leaf blight Downy mildew Neck rot Purple blotch	3	Follow a protective spray schedule starting when diseases are first reported in the areas and repeat at 7 day intervals throughout the season. The addition of a surfactant to spray solutions will improve performance.	Do not apply within 7 days of harvest. Do not apply to exposed bulbs. Do not apply more than 30 lbs (24 lbs active) per acre per crop.
	Smut		Apply 3 lbs per acre as a furrow drench at time of planting onion seeds. Use 75 to 125 gallons of water per acre.	Do not use more than 2.4 lbs active per acre (29,000 linear feet of furrow) with an 18 inch row spacing.
Papaya	Anthracnose (Colletotricum) Phytophthora fruit rot Black spot (Cercospora)	2 to 2.5 (minimum 50 gals per acre)	Begin at flowering; treat central column crown, blossom area and developing fruit. Repeat at 14 to 21 day intervals.	Do not use more than 35 lbs (28 lbs active) per acre per crop. Minimum pre-harvest interval (0 days).
Peanuts	Cercospora leaf spot Rust	1 to 2	Start application when disease first appears or is reported in area. Repeat sprays at 7 to 14 day intervals. Reduce sprays to a 7 day interval during humid weather.	Do not apply within 14 days of harvest. Do not use more than 16 lbs (12.8 lbs active) per acre per crop. Do not feed treated vines to livestock.
Peanuts (Tank mix with "Benlate" Fungicide)	Ascochyta web blotch Cercospora leaf spot Rust	1 1/2 plus 4 ozs "Benlate"	Begin 35 to 40 days after planting or when disease threatens. Repeat at 7 to 14 day intervals (7 to 10 days for rust).	Do not apply within 14 days of harvest. Do not use more than 12 lbs Manzate 200 (9.6 lbs active) per acre per crop. Do not feed treated vines to livestock.
Pears (See Pomefruit)				

CROP	DISEASES CONTROLLED	RATE OF MANZATE 200 PER APPLICATION LBS/ACRE	TIMING/INTERVALS (Also refer to Directions for Use)	RESTRICTIONS/COMMENTS
Pineapple (Seed-Piece Treatment Only)	Phytophthora heart rot	3.2 lbs/10 gals Use up to 100 gals of suspension to treat clean propagation material for one acre.	Dip planting material in fungicide suspension prior to planting. Stir frequently to prevent settling.	Prepare new suspension when 2/3 of volume is used or sooner if noticeably discolored by soil from plant material.
Pomefruits Apples Pears Crabapples Quince	Rusts Scab Fabrea leaf spot	6.0 Maximum per acre use rate based on thorough coverage dilute sprays. Use 2 lbs per 100 gal, up to 300 gal per acre Consult State Extension Service if necessary to adjust for variable tree size.	Pre-Bloom Use: Begin application at 1/4 to 1/2 inch green tip and continue on a 7 to 10 day schedule through bloom. Use either the "Pre-Bloom" or "Extended Application" schedule. DO NOT COMBINE OR INTEGRATE THE TWO TREATMENT SCHEDULES	Do not apply more than 6 lbs (4.8 lbs active) per acre per application. Do not apply after bloom. Do not apply more than 24 lbs (19.2 lbs active) per acre per year. Do not graze livestock in treated areas. It is recommended that this product be used in an Integrated Pest Management Program.
Pomefruits Apples Pears Crabapples Quince	Rusts Scab Fabrea leaf spot	3.0 Maximum per acre use rate based on thorough coverage dilute sprays. Use 1 lb per 100 gal, up to 300 gal per acre Consult State Extension Service if necessary to adjust for variable tree size.	Extended Application Schedule or for Use in Tank Mixtures: For implementation of IPM programs, applications based on tree-row volume, or for use as a resistance management tool; begin applications at 1/4 to 1/2 inch green tip and continue applications on a 7 to 10 day schedule through the second cover spray. Use either the "Pre-Bloom" or "Extended Application" schedule. DO NOT COMBINE OR INTEGRATE THE TWO TREATMENT SCHEDULES	Do not apply more than 3 lbs (2.4 lbs active) per acre per application. Do not apply within 77 days of harvest. Do not apply more than 21 lbs (16.8 lbs active) per acre per year. Do not graze livestock in treated areas. It is recommended that this product be used in an Integrated Pest Management Program.

CROP	DISEASES CONTROLLED	RATE OF MANZATE 200 PER APPLICATION LBS/ACRE	TIMING/INTERVALS (Also refer to Directions for Use)	RESTRICTIONS/COMMENTS
Potatoes	Early blight Late blight	1 to 2	Begin applications when plants are 4 to 6 inches high by applying 1 lb per acre. As the vines increase in size, apply 1 1/2 to 2 lbs per acre at intervals of 5 to 10 days. The addition of a surfactant to spray solutions will improve performance.	Do not apply more than 14 lbs (11.2 lbs active) per acre per crop. Do not use within 3 days of harvest in CT, DE, FL, MA, ME, MI, NH, NY, OH, PA, RI, VT, WI, and within 14 days elsewhere. Vine-kill should occur 14 days before harvest. It is recommended that this product be used in an Integrated Pest Management Program.
Seedpiece (treatment)	Fusarium decay Seedborne common scab	1 1/4 per 50 gal	Dip whole or cut potato tubers in 1 1/4 lbs Manzate 200 fungicide per 50 gallons of water. Place treated tubers in a clean container following treatment and plant as soon as possible. Spread treated seedpieces in a cool place if held before planting.	Do not use treated seed potatoes for food or feed purposes.
Squash (Summer squash)	Downy mildew	2 to 3	Start applications when plants are in the two-leaf stage and repeat at 7 to 10 day intervals. Use sufficient water and direct spray to provide thorough coverage of both upper and lower leaf surfaces.	Do not apply within 5 days of harvest. Do not apply more than 24 lbs (19.2 lbs active) per acre per crop.
Sugar Beets	Cercospora leaf spot	1 1/2 to 2	Begin when disease first threatens. Repeat at 7 to 10 day intervals.	Do not apply within 14 days of harvest. Do not apply more than 14 lbs. (11.2 lbs active) per acre per crop. Do not feed treated sugar beet tops to livestock.
Tomatoes (East of the Mississippi River)	Anthracnose Early blight Gray leaf spot Late blight Leaf mold Septoria leaf spot	3/4 to 1 1/2	Start application when seedlings emerge or transplants are set. Repeat at 3 to 7 day intervals throughout the season. The addition of a surfactant to spray solutions will improve performance.	Do not apply within 5 days of harvest. Do not apply more than 21 lbs (16.8 lbs active) per acre per crop.
		1 1/2 to 3	Start application when seedlings emerge or transplants are set. Repeat at 7 to 10 day intervals throughout the season. The addition of a surfactant to spray solutions will improve performance.	
Tomatoes (West of the Mississippi River)	Anthracnose Early blight Gray leaf spot Late blight Leaf mold Septoria leaf spot	3/4 to 1	Start application when seedlings emerge or transplants are set. Repeat at 3 to 7 day intervals throughout the season. The addition of a surfactant to spray solutions will improve performance.	Do not apply within 5 days of harvest. Do not apply more than 8 lbs (6.4 lbs active) per acre per crop.
		1 1/2 to 2	Start application when seedlings emerge or transplants are set. Repeat at 7 to 10 day intervals throughout the season. The addition of a surfactant to spray solutions will improve performance.	
Watermelon (See Melons)				
Wheat (See Barley)				

TURF GRASSES - Applications restricted to lawn grasses by professional applicators. Not for homeowner use.

CROP	DISEASE/ PEST	RATE	TIMING INTERVAL	LAST APPLICATION TO HARVEST	LIVESTOCK GRAZING/ FEEDING	COMMENTS
Sod Farms	Algae	6 ozs in 3 to 5 gals/1000 sq ft (16 lbs in 130-220 gals/acre).	Begin when algae first appears. Repeat at 7-day intervals as long as condition persists.		Do not graze treated areas or feed clippings to livestock.	Do not use on grasses grown for seed.
	Leaf stem, strip rust	3-4 ozs in 3 to 5 gals/1000 sq ft (8-11 lbs in 130-220 gals/acre).	Begin when disease threatens. Repeat at 7 to 10-day intervals as long as disease persists.		Do not graze treated areas or feed clippings to livestock.	Do not use on grasses intended for grazing, such as range or pasture grasses.
	Leaf spot (Helminthosporium spp.) Rhizoctonia solani brown patch	3-4 ozs in 3 to 5 gals/1000 sq ft (8-11 lbs in 130-220 gals/acre)	Begin when disease threatens.		Do not graze treated areas or feed clippings to livestock.	When conditions are unusually favorable for disease, use 5-8 ozs/1000 sq ft (16-22lbs acre) and reduce intervals to 3 to 5 days.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Seed treatments and professional applications to lawn grasses, golf courses, industrial (office park), municipal and residential lawns are not within the scope of the Worker Protection Standard.

Do not enter treated areas without protective clothing until sprays have dried.

SEED TREATMENTS

LABEL TREATED SEED "Do not use for food, feed or oil purposes. This seed treated with 'Manzate' 200 Fungicide."

For seed treatment, a dye must be added to the treating slurry so that an unnatural color will distinguish the seed as treated.

CROP	DISEASES	SEED TREATMENT RATE -- APPLY AS A SLURRY	
		Ozs/Bu	Ozs/100 lb
Barley	Bunt, Covered smut, Damping-off, Fake loose smut, Seed decay, Seedling blights	1.3 to 2.0	2.7 to 4.2
Corn	Damping-off, seed rot, seedling blight	1.5 to 3.0	2.7 to 5.4
Cotton Acid Delinted	Damping-off, Seedling blights	-	3
Cotton Reginned	Damping-off, Seedling blights	-	6
Flax	Seed decay, Seedling blights, Damping-off	2.0 to 4.0	3.5 to 7.1
Oats	Damping-off, Seedling blights, Seed decay, Smuts	1.3 to 2.0	4.0 to 6.3
Peanuts (shelled)	Damping-off, Seed rots, Seedling blights	2.0 to 4.0	8.0 to 16.0
Rice	Achlya, other soil and seedborne fungi causing seed rot and reduced seedling vigor.	-	2.0 to 4.0 Apply before, during, or after soaking in water.
Rye	Bunt, Covered smut, Damping-off, Seed decay, Seedling blights	1.3 to 2.0	2.3 to 3.6
Safflower	Puccinia carthami (which causes foot-and-rot disease and foliage rust disease)	-	2.0
Sorghum	Covered kernel smut, Damping-off, Seedling blights, Seed rots	1.5 to 2.5	2.7 to 4.5
Tomatoes	Damping-off, Seedling blights, Seed rots	-	8.0
Wheat	Bunt, Covered smut, Damping-off, Seed decay, Seedling blights	1.3 to 2.0	2.2 to 3.3

STORAGE AND DISPOSAL

STORAGE: Important—Never allow “Manzate” 200 to become wet during storage. This may lead to certain chemical changes which will reduce the effectiveness of “Manzate” 200 as a fungicide and create vapors which may be flammable. Store product in original container only. Do not contaminate water, other pesticides, fertilizer, food or feed in storage. Not for use or storage in or around the home. Keep container closed when not in use.

PRODUCT DISPOSAL: Do not contaminate water, food or feed by disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read This Limitation of Warranty and Liability Before Buying or Using This Product. If the Terms Are Not Acceptable, Return the Product at Once, Unopened, and the Purchase Price Will Be Refunded.

It is impossible to eliminate all risks associated with the use of this product. Such risks arise from weather conditions, soil factors, off target movement, unconventional farming techniques, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of DuPont. These risks can cause: ineffectiveness of the product; crop injury, or; injury to non-target crops or plants.

DuPont does not agree to be an insurer of these risks. **WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.**

DuPont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

DUPONT MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

IN NO EVENT SHALL DUPONT OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BUYER'S OR USER'S BARGAINED-FOR EXPECTATION IS CROP PROTECTION. THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF DUPONT OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT OR STRICT LIABILITY), WHETHER FROM FAILURE TO PERFORM OR INJURY TO CROPS OR OTHER PLANTS, AND RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF DUPONT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

DuPont or its Authorized Retailer must have prompt notice of any claim so that an immediate inspection of buyer's or user's growing crops can be made. Buyer and all users shall promptly notify DuPont or a DuPont Authorized Retailer of any claims, whether based on contract, negligence, strict liability, other tort or otherwise or be barred from any remedy.

This Limitation of Warranty and Liability may not be amended by any oral or written agreement.

NET CONTENTS 6 LBS

SL - 288-1 9116 11/1896