

PLEASE NOTE

This image contains more than one label approved for this product on this date.

352-342

9-23-1974



AGRICULTURAL BULLETIN



LANNATE® METHOMYL INSECTICIDE
Oranges, Lemons, Grapefruit
and Tangelos,
California and Arizona

SUPPLEMENTAL LABELING
EPA REG. NO. 352-342

LANNATE® METHOMYL INSECTICIDE
FOR USE ON ORANGES, LEMONS, GRAPEFRUIT, AND TANGELOS
IN CALIFORNIA AND ARIZONA

Du Pont "Lannate" Methomyl Insecticide is recommended for use in California and Arizona on oranges, lemons, grapefruit, and tangelos for control of citrus thrips.

HOW TO USE

Apply "Lannate" at 1/2 to 1 lb. per acre. Use sufficient water to obtain thorough coverage (use a minimum of 15 gals./acre by air). Use the higher rate on severe infestations of thrips. Apply in the early spring before bloom when the new growth is about 3 to 4 inches long. Make additional applications as needed until the new fruit is walnut size. Application at petal fall may be critical to prevent fruit scarring. Applications during mid-summer to protect new growth on young trees are also recommended.


Note: Do not apply within 1 day of harvest. Do not graze livestock in treated orchards for 10 days after treatment. This product is toxic to bees and should not be applied when bees are actively visiting the area.

IMPORTANT

BEFORE USING "LANNATE", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental directions for use of this product which may not appear on package label. Follow the instructions carefully.

2/19/74-A



NEXT

LABEL



352-342

9-23-1974

AGRICULTURAL BULLETIN



LANNATE® METHOMYL INSECTICIDE
California and Arizona
SUGAR BEETS'

1/1

ACCEPTED
SEP 23 1974
UNDER THE FEDERAL
FUNGICIDE AND WEED
KILLING ACT
FOR RECORDED
ED UNDER NO. 352-342

SUPPLEMENTAL LABELING
EPA REG. NO. 352-342

LANNATE® METHOMYL INSECTICIDE
FOR USE ON SUGAR BEETS IN
CALIFORNIA AND ARIZONA

Du Pont "Lannate" Methomyl Insecticide is recommended for use on sugar beets for the control of beet armyworm, beet webworm, flea beetle and aphids in California and Arizona.

HOW TO USE

Apply 1/4 to 1 lb. "Lannate" per acre in sufficient water to obtain thorough coverage. Make the first application when insects appear and plant damage occurs. Make 2 or 3 additional applications if needed to maintain control.

For effective insect control, proper spray timing and coverage are essential. Best control is obtained when spray schedules are initiated on small insects. In general, use the low rate on light insect infestations. In beet armyworm control, use the dosage rates below 1/2 lb. "Lannate" per acre only when treating small plants.

Note: Do not apply within 7 days of harvest. Do not feed treated tops to livestock.

IMPORTANT

BEFORE USING "LANNATE", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental directions for use of this product, which may not appear on package label. Follow the instructions carefully.

1/25/74

NEXT

LABEL

352-342

09/23/1974



1/1



AGRICULTURAL BULLETIN

LANNATE® METHOMYL INSECTICIDE
NECTARINES

SUPPLEMENTAL LABELING
EPA REG. NO. 352-342

ACCEPTED

SEP 23 1974

SMITHSONIAN INSTITUTION
FEDERAL BUREAU OF INVESTIGATION
FOR THE U.S. DEPARTMENT OF JUSTICE
RECEIVED
SEP 23 1974
EPA REG. NO. 352-342

LANNATE® METHOMYL INSECTICIDE
FOR USE ON NECTARINES
IN THE STATES OF CALIFORNIA AND ARIZONA

Du Pont "Lannate" Methomyl Insecticide is recommended for use on nectarines in California and Arizona for control of thrips.

HOW TO USE

Apply 1/2 to 1 lb. per acre. Use sufficient water (use a minimum of 10 gals. per acre by air) to obtain thorough coverage.

Make first application at early bloom when thrips are present in damaging numbers. Make one or two additional applications if needed to maintain control.


Note: "Lannate" is toxic to bees and should not be applied when bees are actively visiting the area. Do not treat within 1 day of harvest. Do not graze livestock in treated orchards for 10 days after treatment. Pilot should not assist in the mixing and loading operation.

IMPORTANT

BEFORE USING "LANNATE", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental directions for use of this product which may not appear on package label. Follow the instructions carefully.

2/4/74-A



NEXT

LABEL



352-342

09/23/1974

AGRICULTURAL BULLETIN




1/1

LANNATE® METHOMYL INSECTICIDE
Apples
East of Rocky Mountains

SUPPLEMENTAL LABELING
EPA REG. NO. 352-342

LANNATE® METHOMYL INSECTICIDE
FOR USE ON APPLES IN AREAS
EAST OF THE ROCKY MOUNTAINS



Du Pont "Lannate" Methomyl Insecticide is recommended for use on apples to control certain aphids, codling moth, and red-banded leafroller in areas east of the Rocky Mountains.

HOW TO USE

Apple Aphid, Rosy Apple Aphid - Apply ^{0.20} 1/4 to ^{0.45} 1/2 lb. per 100 gals. to established infestations.

Codling Moth - Make 4 to 7 applications of ^{0.20} 1/4 to ^{0.45} 1/2 lb. per 100 gals. at 10- to 12-day intervals beginning at the second cover spray.

Red-Banded Leafroller (Northeast and Central U.S. only) - Apply 1/2 lb. per 100 gals. at green tip to 1/2 inch green and at petal fall and first cover.

Apply in sufficient water to obtain thorough coverage (run-off) but not in excess of 400 gals. per acre. For concentrate sprayers use proportionately less water.

Spray Application - Fill spray tank 1/4 to 1/2 full of water. Add "Lannate" directly to spray tank. Mix thoroughly to fully dissolve the insecticide; once dissolved continued agitation is not required. Use mechanical or hydraulic means; do not use air agitation.


Do not apply within 28 days of harvest. Do not graze livestock in treated orchards for 10 days after treatment.

IMPORTANT

BEFORE USING "LANNATE", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental directions for use of this product which may not appear on package label. Follow the instructions carefully.

2/19/74-A



data submitted 2/20/74; accepted 5/30/74 THH conditionally, revised label accepted

NEXT

LABEL

352-342

09/23/1974



1/1

AGRICULTURAL BULLETIN

LANNATE[®] METHOMYL INSECTICIDE
Cotton
East of Rocky Mountains

ACCEPTED
SEP 23 1974
352-342

SUPPLEMENTAL LABELING
EPA REG. NO. 352-342

LANNATE[®] METHOMYL INSECTICIDE
FOR LATE SEASON USE ON COTTON
EAST OF THE ROCKY MOUNTAINS

Du Pont "Lannate" Methomyl Insecticide is recommended for late season use on cotton east of the Rocky Mountains to control bollworm and tobacco budworm.

HOW TO USE

Apply 1/2 to 3/4 lb. "Lannate" per acre. Make the first application after the desired crop (boll load) is set on the plant. Repeat at 3 to 5 day intervals as needed. Do not apply more than three treatments. Apply "Lannate" by conventional ground or aerial application equipment. Use sufficient water to obtain thorough coverage (use a minimum of 3 gals. per acre for aerial application). Do not use surfactants in the spray.

"Lannate" may be applied up to 15 days before harvest. Do not graze or feed treated cotton to livestock.

NOTE: USE OF "LANNATE" MAY REDDEN COTTON FOLIAGE. IF REDDENING IS EXCESSIVE, DISCONTINUE USE OF "LANNATE" OR ALTERNATE WITH OTHER INSECTICIDES.

IMPORTANT

BEFORE USING "LANNATE", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

2/5/74-A

NEXT

LABEL

352-342
**AGRICULTURAL
BULLETIN**

09/23/1974

1/1

LANNATE® METHOMYL INSECTICIDE
California and Arizona
SUGAR BEETS,

ACCEPTED
9-23-74
SEP 23 1974

SUPPLEMENTAL LABELING
EPA REG. NO. 352-342

LANNATE® METHOMYL INSECTICIDE
FOR USE ON SUGAR BEETS IN
CALIFORNIA AND ARIZONA

352-342

Du Pont "Lannate" Methomyl Insecticide is recommended for use on sugar beets for the control of beet armyworm, beet webworm, flea beetle and aphids in California and Arizona.

HOW TO USE

Apply 1/4 to 1 lb. "Lannate" per acre in sufficient water to obtain thorough coverage. Make the first application when insects appear and plant damage occurs. Make 2 or 3 additional applications if needed to maintain control.

For effective insect control, proper spray timing and coverage are essential. Best control is obtained when spray schedules are initiated on small insects. In general, use the low rate on light insect infestations. In beet armyworm control, use the dosage rates below 1/2 lb. "Lannate" per acre only when treating small plants.

Note: Do not apply within 7 days of harvest. Do not feed treated tops to livestock.

IMPORTANT

BEFORE USING "LANNATE", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental directions for use of this product which may not appear on package label. Follow the instructions carefully.

1/25/74

application 7/15/74 to make changes: geographical restriction to Ariz. & Calif. + above, accepted 9/23/74

NEXT

LABEL




AGRICULTURAL BULLETIN



LANNATE® METHOMYL INSECTICIDE
CELERY

SUPPLEMENTAL LABELING
EPA REG. NO. 352-342

LANNATE® METHOMYL INSECTICIDE
FOR USE ON CELERY



Du Pont "Lannate" Methomyl Insecticide is recommended for use on celery for control of cabbage looper and beet armyworm.

HOW TO USE

Apply "Lannate" at 5- to 7-day intervals or as needed in sufficient water (5-15 gal. per acre by air) to obtain uniform coverage.

- Beet Armyworm - apply 1/2 to 1 lb. per acre.
- Cabbage looper - apply 1 lb. per acre.

Do not apply within 14 days of harvest. *Apply before 7/31/74*

Note: Pilot should not assist in the mixing and loading operation.

IMPORTANT

BEFORE USING "LANNATE", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental directions for use of this product which may not appear on package label. Follow the instructions carefully.

6/14/74

