

PLEASE NOTE

This image contains more than one label approved for this product on this date.

352-342

03/26/1975

1/1



AGRICULTURAL BULLETIN



LANNATE® METHOMYL INSECTICIDE
DRY BEANS

SUPPLEMENTAL LABELING
EPA REG. NO. 352-342

LANNATE® METHOMYL INSECTICIDE
FOR USE ON DRY BEANS



Du Pont "Lannate" Methomyl Insecticide is recommended for use on dry beans for control of leafhoppers.

HOW TO USE

Apply 1/4 to 1 lb. "Lannate" per acre in sufficient water (5 to 15 gals. per acre by air) to obtain uniform coverage. Make application when insect first appears and repeat at 5- to 7-day intervals or as needed. Make no more than 2 applications after pod set.


NOTE: Do not harvest dry beans within 25 days of last application. Vines may not be foraged within 3 days of application or cut for hay within 7 days of application.

IMPORTANT

BEFORE USING "LANNATE", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

12/12/74



data submitted 7/15/74; accepted 11/13/74 POH on OBJ, revised label accepted

NEXT

LABEL

352-342

3/26/1975



AGRICULTURAL BULLETIN



LANNATE® METHOMYL INSECTICIDE
ONIONS
STATE OF ARIZONA

SUPPLEMENTAL LABELING
EPA REG. NO. 352-342



LANNATE® METHOMYL INSECTICIDE
FOR USE ON ONIONS
IN THE STATE OF ARIZONA

Du Pont "Lannate" Methomyl Insecticide is recommended for use on onions (green or dry) for control of beet armyworm and thrips in the State of Arizona.

HOW TO USE

Apply 1/2 lb. "Lannate" per acre in sufficient water (5 to 15 gals. per acre by air) to obtain uniform coverage. Apply when insect first appears. Make one additional application if needed. Add a wetting agent.


NOTE: Do not apply within 28 days of harvest on green onions or 7 days of harvest on dry onions.

IMPORTANT

BEFORE USING "LANNATE", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

12/12/74



data submitted 7/26/74; accepted 11/13/74 POH on OBJ, revised label accepted

NEXT

LABEL

352-342

3-26-1975



AGRICULTURAL BULLETIN



LANNATE® METHOMYL INSECTICIDE
CARROTS
ARIZONA AND CALIFORNIA

SUPPLEMENTAL LABELING
EPA REG. NO. 352-342

LANNATE® METHOMYL INSECTICIDE
FOR USE ON CARROTS
IN THE STATES OF ARIZONA AND CALIFORNIA



Du Pont "Lannate" Methomyl Insecticide is recommended for use on carrots to control beet armyworm in the states of Arizona and California.

HOW TO USE

Apply 1/3 to 1 lb. "Lannate" per acre in sufficient water (5 to 15 gals. per acre by air) to obtain uniform coverage. Make first application when insects first appear and repeat at 5- to 7-day intervals or as needed.


NOTE: Do not apply within 1 day of harvest

IMPORTANT

BEFORE USING "LANNATE", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

12/12/74



data submitted 12/19/74; accepted 3/26/75 POH

0.2 ppm root reg. abax FR 2/26 7

NEXT

LABEL

352-342

3-26-1975



AGRICULTURAL BULLETIN



LANNATE® METHOMYL INSECTICIDE
BLACKEYED PEAS
STATE OF CALIFORNIA

SUPPLEMENTAL LABELING
EPA REG. NO. 352-342

LANNATE® METHOMYL INSECTICIDE
FOR USE ON BLACKEYED PEAS
IN THE STATE OF CALIFORNIA



Du Pont "Lannate" Methomyl Insecticide is recommended for use on blackeyed peas for control of leafhoppers, beet armyworm, and cabbage looper in the State of California.

HOW TO USE

Apply "Lannate" in sufficient water (5 to 15 gals. per acre by air) to obtain uniform coverage. Make application when insect first appears and repeat at 5- to 7-day intervals or as needed. Make no more than 2 applications after pod set.

Leafhoppers - apply 1/4 to 1 lb. per acre.
Beet armyworm, cabbage looper - apply 1/2 to 1 lb. per acre.


NOTE: Do not harvest blackeyed peas within 25 days of last application. Vines may not be foraged within 3 days of application or cut for hay within 7 days of application.

IMPORTANT

BEFORE USING "LANNATE", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

12/12/74



data submitted 7/15/74; accepted 11/13/74 FOI on OBI, revised label accepted