

PLEASE NOTE

This image contains more than one label approved for this product on this date.

352-342


08/18/1975




AGRICULTURAL BULLETIN



LANNATE® METHOMYL INSECTICIDE
Grapes
California and Arizona



SUPPLEMENTAL LABELING
EPA REG. NO. 352-342



LANNATE® METHOMYL INSECTICIDE
FOR USE ON GRAPES IN
CALIFORNIA AND ARIZONA

Du Pont "Lannate" Methomyl Insecticide is recommended for use on grapes for control of leafhoppers in California and Arizona.

HOW TO USE

Apply 1/2 to 1 lb. per acre by ground equipment in sufficient water to obtain thorough coverage.

Make the first application when insects appear in damaging numbers and repeat at 5- to 7-day intervals or as needed. Do not make more than 2 applications after fruit set.

Note: Do not treat within 1 day of harvest; do not treat grapes for wine within 14 days of harvest.

IMPORTANT

BEFORE USING "LANNATE", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

10/4/73-B*

NEXT

LABEL



AGRICULTURAL BULLETIN



1/2

LANNATE® METHOMYL INSECTICIDE

Grapes

West of the Rocky Mountains

ACCEPTED
AUG 18 1975
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 352-342

SUPPLEMENTAL LABELING
EPA REG. NO. 352-342

LANNATE® METHOMYL INSECTICIDE
FOR USE ON GRAPES
WEST OF THE ROCKY MOUNTAINS

Du Pont "Lannate" Methomyl Insecticide is recommended for use on grapes west of the Rocky Mountains for control of omnivorous leafroller, grape leaffolder, orange tortrix and salt marsh caterpillar.

HOW TO USE

Apply by ground equipment in sufficient water to obtain thorough coverage. Good coverage is particularly important in the case of insects which directly attack the grape berries.

Omnivorous Leafroller - Apply 1 lb. per acre. Make the first application when insects appear in damaging numbers or when leaf folding is observed. Repeat at 5- to 7-day intervals or as needed.

Grape Leaffolder - Apply 1 lb. per acre. Make the first application when insects appear in damaging numbers or when leaf rolling is observed. Repeat at 5- to 7-day intervals or as needed.

Orange Tortrix - Apply 1 lb. per acre. Make the first application when insects appear in damaging numbers in spring growth. Repeat at 5- to 7-day intervals or as needed.

Salt Marsh Caterpillar - Apply 1/2 to 1 lb. per acre. Make the first application when insects appear in damaging numbers. For aerial application, use a minimum of 10 gals. of spray mixture per acre.

Note: Do not treat within 1 day of harvest; do not treat grapes for wine within 14 days of harvest.



