CROPS	Method of application: A means Air; G means Ground	INSECTS	LBS. PER ACRE	LAST APPLICATION—DAYS To Harvest	LAST APPLICATION—DAYS To Livestock Grazing/Feeding
Celery	A, G	Cabbage Looper	1	14	
		Earworm Whorl as needed	1/2-1/2		
Corn (Sweet)		Earworm Ears [illegible] days or as needed	1/4-1/2	0	3
		Fall Armyworm European Corn Borer		Ear	Fodder
	A, G	Ears [illegible] days as needed	1/2		
Cucumber	G	Cabbage Looper	1/2-1	3	
Lettuce (Head)		Cabbage and Alfalfa Loopers	1/4-1/2	7	
	A, G	Beet Armyworm	1/2-1	10	
Melons	G	Cabbage Looper	1/2-1	3	
Peanuts—East of Miss. River	G	Corn Earworm Up to [illegible] applications as needed	1/4-1	21	Do not feed treated vines
Peppers	A, G	Green Peach Aphid	1/2	10	
		Tuberworm Cabbage Looper	1/2		
Potato		Aphids	1/2-1	14	
	A, G	Leafhoppers, Flea Beetles } East of Miss. River	1/2		
Soybeans—(Southeastern United States)	G	Soybean Looper Green Cloverworm Mexican Bean Beetle Corn Earworm Up to 2 applications	1/4-1/2	14	3 (forage) 7 (hay)
Spinach	G	Cabbage Looper Beet Armyworm Do not apply when min. daily temp is 32° F or lower Do not apply to seedlings less than 3" in diameter	1/2-1	16	
Squash (Summer)	G	Cabbage Looper Melonworm, Pickleworm } Southeast only	1/2-1	3	
Tomato		Tomato Fruitworm Aphids	1/2	1	
	A, G	Cabbage Looper Beet Armyworm	over 1/2-1	2	
Tobacco (Except shade)		Flea Beetle Hornworm	1/4-1/2	7 (flue cured)	
		Budworm (2-3 applications before flower buds open) Cabbage Looper	1/2	14 (air or fire cured)	
	A, G	Aphids			
Chrysanthemums	G	Cabbage Looper Corn Earworm Beet Armyworm Thrips weekly	1/4-1/2 lb. 100 gals		

DU PONT Lannate® NET 2 LBS.

METHOMYL INSECTICIDE — **WATER SOLUBLE POWDER**

ACTIVE INGREDIENT: [illegible] **90%**

INERT INGREDIENTS **10%**

EPA Reg. No 352-342 AA

U.S. Pats. 3,576,834 & 3,639,633

Keep out of reach of children.

DANGER POISON

See Antidote and Precautions in Right Panel. ▶

MAY BE FATAL IF SWALLOWED. POISONOUS IF INHALED

Do not breathe dust or spray mist. Do not get in eyes, on skin, on clothing.

In case of contact, wash skin with plenty of soap and water; for eyes, flush with water for 15 minutes and get medical attention. Remove and wash contaminated clothing before re-use.

IMPORTANT—When mixing, spraying, loading, applying or otherwise handling wear goggles and a mask or respirator of a type suitable for protection against insecticide dust or spray mist [illegible] Methomyl dust. Agri-Tox R5 [illegible] Wash clothes daily, [illegible] Wash clean clothes daily. [illegible] contaminated [illegible] and washing [illegible]

[illegible]

E. I. DU PONT DE NEMOURS & COMPANY (INC.)
BIOCHEMICALS DEPARTMENT WILMINGTON DELAWARE

POISON

ANTIDOTE AND FIRST AID

ATROPINE IS AN ANTIDOTE—CONSULT PHYSICIAN FOR EMERGENCY SUPPLY OF 1/100 GRAIN ATROPINE TABLETS. CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED POISONING.

If swallowed, give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. **If inhaled,** remove from exposure. Have patient lie down and keep quiet. If patient is not breathing, start artificial respiration immediately. **Never give anything by mouth to an unconscious person.**

If warning symptoms appear (See **warning symptoms** below), before physician arrives, immediately swallow two atropine tablets (each 1/100 gr.); thereafter, every 10 to 15 minutes, take one atropine tablet (1/100 gr.) until throat becomes dry and skin becomes dry and flushed. Take additional tablets as necessary to maintain a moderately dry throat and dry, flushed skin until physician arrives.

NOTE TO PHYSICIAN

WARNING SYMPTOMS—Methomyl poisoning produces effects associated with anticholinesterase activity which may include weakness, blurred vision, headache, nausea, **abdominal cramps, discomfort in the chest, constriction of pupils, sweating, slow pulse, muscle tremors.**

TREATMENT—Atropine sulfate should be used for treatment. Administer repeated doses, 1.2 to 2.0 mg. intravenously every 10 to 30 minutes until full atropinization is achieved. Maintain atropinization until the patient recovers. Artificial respiration or oxygen may be necessary. Allow no further exposure to any cholinesterase inhibitor until recovery is assured.

Do not use morphine or 2-PAM.

NOTICE TO BUYER—Seller warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. This warranty does not extend to use of this product contrary to label use directions, or under abnormal use conditions, or under conditions not reasonably foreseeable to seller; buyer assumes all risk of any such use. Seller makes no other warranties, express or implied.

Purchase of this material does not confer any rights under patents of countries outside of the United States

CONTINUED ON RIGHT PANEL ▶