

352-344

01/27/1976

1/1




AGRICULTURAL BULLETIN



LANNATE® METHOMYL INSECTICIDE
COTTON
EAST OF THE ROCKY MOUNTAINS

SUPPLEMENTAL LABELING
EPA REG. NO. 352-342



LANNATE® METHOMYL INSECTICIDE
FOR USE ON COTTON
EAST OF THE ROCKY MOUNTAINS

Du Pont "Lannate" Methomyl Insecticide is recommended for occasional use with other insecticides on cotton east of the Rocky Mountains for control of bollworm, tobacco budworm and cotton leafperforator.

HOW TO USE

Apply 1/2 lb. "Lannate" per acre. Use as an occasional treatment in a regular spray schedule. Do not use more often than every 10 days. Apply in sufficient water (minimum 3 gals. per acre by air) to obtain thorough coverage.

Note: Use of "Lannate" may redden cotton foliage. If reddening is excessive, discontinue use of "Lannate" or alternate with other insecticides. "Lannate" may be applied up to 15 days before harvest. Do not graze or feed treated cotton to livestock.

IMPORTANT

BEFORE USING "LANNATE", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

10/29/75