

PLEASE NOTE

This image contains more than one label approved for this product on this date.

352-342


05/27/1976



AGRICULTURAL BULLETIN

LANNATE[®] METHOMYL INSECTICIDE
TOBACCO

SUPPLEMENTAL LABELING
EPA REG. NO. 352-342



LANNATE[®] METHOMYL INSECTICIDE
FOR USE ON TOBACCO (EXCEPT SHADE TOBACCO)

Du Pont "Lannate" Methomyl Insecticide is recommended for control of flea beetle, tobacco hornworm, cabbage looper, tobacco budworm and aphids.

HOW TO USE

Apply as directed on the package label.


Note: Do not apply within 5 days of harvest on tobacco to be flue cured. Do not apply within 14 days of harvest on tobacco to be air or fire cured. The maximum use rate is 1/2 lb. "Lannate" per acre per treatment.

IMPORTANT

BEFORE USING "LANNATE", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

3/5/76



NEXT

LABEL

352-342

5-27-1976




1/1



AGRICULTURAL BULLETIN

LANNATE® METHOMYL INSECTICIDE
California and Arizona
Sugar Beets

SUPPLEMENTAL LABELING
EPA REG. NO. 352-342



LANNATE® METHOMYL INSECTICIDE
FOR USE ON SUGAR BEETS IN
CALIFORNIA AND ARIZONA

Du Pont "Lannate" Methomyl Insecticide is recommended for use on sugar beets for the control of beet armyworm, beet webworm, flea beetle and aphids in California and Arizona.

HOW TO USE

Apply 1/4 to 1 lb. "Lannate" per acre in sufficient water to obtain thorough coverage. Make the first application when insects appear and plant damage occurs. Make 2 or 3 additional applications if needed to maintain control.

For effective insect control, proper spray timing and coverage are essential. Best control is obtained when spray schedules are initiated on small insects. In general, use the low rate on light insect infestations. In beet armyworm control, use the dosage rates below 1/2 lb. "Lannate" per acre only when treating small plants.

Note: Do not apply within 7 days of harvest. Do not feed treated tops to livestock within 30 days of last application.

IMPORTANT

BEFORE USING "LANNATE", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

3/22/76

