

## **PLEASE NOTE**

**This image contains more than one label approved for this product on this date.**

352-342

07/29/1977



# AGRICULTURAL BULLETIN



LANNATE® METHOMYL INSECTICIDE  
SOYBEANS

SUPPLEMENTAL LABELING  
EPA REG. NO. 352-342

CCC  
JUL 29 1977  
Under the Federal Insecticide, Fungicide, and Rodenticide Act, recommended for use pesticide registered under EPA Reg. No. 352-342

LANNATE® METHOMYL INSECTICIDE  
FOR USE ON SOYBEANS

Du Pont "Lannate" Methomyl Insecticide is recommended for use on soybeans for the control of soybean looper.

### HOW TO USE

Apply 1/2 to 1 lb. "Lannate" per acre when insects appear in damaging numbers. Make additional applications as needed to maintain control. Use sufficient water to obtain thorough coverage. Use a minimum of 2 gals. per acre for aerial application.


Note: Do not apply within 14 days of bean harvest. Do not graze or feed treated soybean forage to livestock within 10 days of last application nor soybean hay within 12 days of last application.

### IMPORTANT

BEFORE USING "LANNATE", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

6/6/77-A



**NEXT**

**LABEL**

352-342


7-29-1977



# AGRICULTURAL BULLETIN



"LANNATE" METHOMYL INSECTICIDE  
SOYBEANS



SUPPLEMENTAL LABELING  
EPA REG. NO. 352-342

## "LANNATE" METHOMYL INSECTICIDE FOR USE ON SOYBEANS

Du Pont "Lannate" Methomyl Insecticide is recommended for insect control on soybeans. Low use rates will control light to moderate infestations of velvetbean caterpillar and help conserve beneficial predators.

### HOW TO USE

Apply 1/8 to 1/4 pound per acre before insect populations build to economic levels. Use the lower rate on light infestations. Make up to 3 additional applications as needed to retain control. Use sufficient water to obtain thorough coverage.

Periodic scouting of fields is required to insure proper timing of spray.

Note: Do not apply within 14 days of bean harvest. Do not graze or feed treated soybean forage to livestock within 3 days of last application nor soybean hay within 7 days of last application.

### IMPORTANT

BEFORE USING "LANNATE", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

6/22/77