



AGRICULTURAL BULLETIN



LANNATE® METHOMYL INSECTICIDE
COTTON

ACCEPTED
AUG 15 1977
Under the Federal Insecticide, Fungicide, and Plant Growth Regulator Act, as amended, for the pesticide LANNATE® METHOMYL INSECTICIDE, EPA Reg. No. 352-342

SUPPLEMENTAL LABELING
EPA REG. NO. 352-342

LANNATE® METHOMYL INSECTICIDE
COMBINATION WITH METHYL PARATHION
FOR USE ON COTTON
IN THE MIDSOUTH AND SOUTHEAST

Du Pont "Lannate" Methomyl Insecticide is recommended in combination with methyl parathion for use on cotton for control of bollworm and bollweevil in the Midsouth and Southeast.

HOW TO USE

Apply 1/4 - 1/3 lb. "Lannate" plus 1 - 2 pts. methyl parathion (4 lb./gal. emulsifiable concentrate) in sufficient water (minimum of 2 gals. per acre by air) to obtain thorough coverage. Repeat at 5- to 7-day intervals.

Apply the higher use rates and shorten spray intervals with increasing insect populations.

Note: Use of "Lannate" in combination with methyl parathion may redden cotton foliage. If reddening is excessive, discontinue use of this combination or alternate with other insecticides. "Lannate" + methyl parathion may be applied up to 15 days before harvest. Do not graze or feed treated cotton to livestock. Do not permit workers to handle treated crops within 48 hours after application. Do not hand pick within 7 days of application.

IMPORTANT

BEFORE USING "LANNATE", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

