



AGRICULTURAL BULLETIN



LANNATE® METHOMYL INSECTICIDE
CUCUMBER & MELONS

ACCEPTED
NOV 15 1977
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 352-342

SUPPLEMENTAL LABELING
EPA REG. NO. 352-342

LANNATE® METHOMYL INSECTICIDE
FOR USE ON CUCUMBER AND MELONS IN THE SOUTHEAST

Du Pont "Lannate" Methomyl Insecticide is recommended for use on cucumber and melons to control melonworm, and pickleworm in the southeastern region of the United States.

HOW TO USE

Apply 1/2 to 1 lb. "Lannate" per acre in sufficient water (aerial: 5 to 15 gals. per acre) to obtain thorough coverage. Treat when worms first appear in blossoms or on foliage. Retreat as required to maintain control.

Note: "Lannate" sprays are toxic to bees and should not be applied when bees are actively visiting the area. Do not apply 1/2 lb. within 1 day of harvest. Do not apply over 1/2 lb. within 3 days of harvest.

IMPORTANT

BEFORE USING "LANNATE", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

