

PLEASE NOTE

This image contains more than one label approved for this product on this date.

352-342

01/17/1978

1/1



AGRICULTURAL BULLETIN



LANNATE® METHOMYL INSECTICIDE
SORGHUM *

ACCEPTED
JAN 17 1978
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. **352-342**

SUPPLEMENTAL LABELING
EPA REG. NO. 352-342

LANNATE® METHOMYL INSECTICIDE
FOR USE ON SORGHUM

Du Pont "Lannate" Methomyl Insecticide is recommended for use on sorghum for control of sorghum midge, sorghum webworm and fall armyworm.

HOW TO USE

Sorghum Midge - Use 1/4 to 1/2 lb. "Lannate" per acre. Apply when 50% of the plants have begun to bloom. If adults are active 3 to 5 days after treatment, make additional treatments as needed.

Sorghum Webworm - Apply 1/2 lb. "Lannate" per acre when larvae are found in sorghum heads. Make additional treatments as needed.

Fall Armyworm (budworm) - Apply 1/2 lb. "Lannate" per acre when larvae are found in the whorl. Make additional treatments as needed.

Mix "Lannate" with sufficient water (minimum 3 gals. per acre by air) to obtain uniform coverage.

Note: Do not apply within 14 days of harvest. Do not graze or feed treated plants to livestock within 14 days after application.

IMPORTANT

BEFORE USING "LANNATE", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

1/26/77 -A

NEXT

LABEL

352-342

1-17-1978

1/1



AGRICULTURAL BULLETIN



LANNATE® INSECTICIDE
ASPARAGUS

SUPPLEMENTAL LABELING
EPA REG. NO. 352-342

LANNATE® INSECTICIDE
FOR USE ON ASPARAGUS

ACCEPTED
JAN 17 1978
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 352-342

Du Pont "Lannate" Insecticide is recommended for use on asparagus for control of the asparagus beetle and spotted asparagus beetle.

HOW TO USE

Apply 1/2 to 1 lb. "Lannate" per acre in sufficient water (10-15 gals. per acre by air) to obtain thorough coverage. Apply when insects first appear in damaging numbers and repeat at 5- to 7-day intervals or as needed. Use the highest recommended rate for controlling severe infestation.

Note: Do not apply within 1 day of harvest.

IMPORTANT

BEFORE USING "LANNATE", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

9/12/77

