

**DRUM CONTAINS LANNATE® IN A WATER SOLUBLE BAG —
ENTIRE CONTENTS DISSOLVE IN WATER.**

After opening drum, immediately dump entire contents into spray tank. Do not touch bag; handling may cause breakage.



IMPORTANT—When mixing spray, loading, applying or otherwise handling, wear goggles and a mask or respirator of a type suitable for protection against insecticide dust or spray mist such as: Willson Products "Agri-Tox" R533 Felt 11-P Cartridge or Mine Safety Appliances type H "Comfo". Wear clean clothes daily. Wash thoroughly after handling and before eating or smoking. Pilot should not assist in the mixing and loading operation. Do not contaminate or store near feed or foodstuffs.

Not for use or storage in or around the home. Keep animals and unprotected persons out of operational areas during treatment and while there is danger of drift. Do not re-use container; destroy when empty.

This product is toxic to fish, birds and other wildlife. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is toxic to bees and should not be applied when bees are actively visiting the area. Apply late in evening or early morning where honey bees visit fields. In alfalfa frequented by alkali or leafcutting bees, do not apply during bloom or when these bees will be present in fields.

ACCEPTED
DEC 14 1978
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under Reg. No. 352-342

NOTICE OF WARRANTY
Du Pont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Du Pont. In no case shall Du Pont be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. DU PONT MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.



Water Soluble Powder

ACTIVE INGREDIENT: Methomyl S-methyl N-[(methylcarbamoyloxy)thioacetimidate] 90%
INERT INGREDIENTS 10%
U. S. Pats. 3,576,834 & 3,639,633 EPA Reg. No. 352-342-AA
EPA Est. 352-TX-1

Keep out of reach of children.

DANGER POISON

See Antidote in Right Panel ▶

MAY BE FATAL IF SWALLOWED. POISONOUS IF INHALED.

Do not breathe dust or spray mist.
Do not get in eyes, on skin, on clothing.

In case of contact, wash skin with plenty of soap and water; for eyes, flush with water for 15 minutes and get medical attention; remove and wash contaminated clothing before re-use.

NET 10 LBS.

HANDLE THIS PACKAGE CAREFULLY TO PREVENT BREAKAGE OF INNER BAG WHEN STORED AT LOW TEMPERATURES. ALLOW TO WARM ABOVE 50°F. FOR NORMAL HANDLING.

E. I. du Pont de Nemours & Co. (Inc.)
BIOCHEMICALS DEPARTMENT, WILMINGTON, DELAWARE



ATROPINE IS AN ANTIDOTE—CONSULT PHYSICIAN FOR EMERGENCY SUPPLY OF 1/100 GRAIN ATROPINE TABLETS. CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED POISONING.

If swallowed, drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger or blunt object. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

If inhaled, remove from exposure. Have patient lie down and keep quiet. If patient is not breathing, start artificial respiration immediately. Never give anything by mouth to an unconscious person.

If warning symptoms appear (See warning symptoms below), before physician arrives, immediately swallow two atropine tablets (each 1/100 gr.); thereafter, every 10 to 15 minutes, take one atropine tablet (1/100 gr.) until throat becomes dry and skin becomes dry and flushed. Take additional tablets as necessary to maintain a moderately dry throat and dry, flushed skin until physician arrives.

NOTE TO PHYSICIAN

WARNING SYMPTOMS—Methomyl poisoning produces effects associated with anticholinesterase activity which may include weakness, blurred vision, headache, nausea, abdominal cramps, discomfort in the chest, constriction of pupils, sweating, slow pulse, muscle tremors.

TREATMENT—Atropine sulfate should be used for treatment. Administer repeated doses, 1.2 to 2.0 mg. intravenously every 10 to 30 minutes until full atropinization is achieved. Maintain atropinization until the patient recovers. Artificial respiration or oxygen may be necessary. Allow no further exposure to any cholinesterase inhibitor until recovery is assured.

Do not use morphine or 2-PAM. However, for exposure to "Lannate" and organophosphorous insecticides, 2-PAM may be used as required to supplement the atropine sulfate treatment as described above.

NOTICE TO BUYER—Purchase of this material does not confer any rights under patents of countries outside of the United States.

B-21770 5-78 Made in U.S.A. Printed in U.S.A.

352-342

12/14/1978

1/2

