

PLEASE NOTE

This image contains more than one label approved for this product on this date.

352-342

02/14/1979




AGRICULTURAL BULLETIN



LANNATE® INSECTICIDE
ALFALFA

SUPPLEMENTAL LABELING
EPA REG. NO. 352-342

LANNATE® INSECTICIDE
FOR USE ON ALFALFA



Du Pont Lannate® Insecticide is recommended for use on alfalfa for the control of pea aphid. Treatment often results in a favorable ratio of predators to prey.

HOW TO USE

Apply 1/2 to 1 lb Lannate® per acre in sufficient water (minimum 5 gals per acre by air) to obtain thorough coverage of plants. Apply when insects appear in damaging numbers. Repeat application to maintain control.

- Note:
- 1) Do not apply to dormant or semi-dormant alfalfa when minimum daily temperature is 50°F or lower.
 - 2) Do not graze or feed treated alfalfa to livestock within 7 days of last application.
 - 3) This product is toxic to bees and should not be applied when bees are actively visiting the area. In fields visited by honeybees, make application late in the evening or early morning. In areas frequented by alkali or leafcutting bees, do not apply during bloom or when the bees will be present in fields.

IMPORTANT

BEFORE USING LANNATE®, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

9/19/78

NEXT

LABEL

352-342

2/14/1979

1/1



AGRICULTURAL BULLETIN



LANNATE® INSECTICIDE
SWEET CORN

SUPPLEMENTAL LABELING
EPA REG. NO. 352-342

LANNATE® INSECTICIDE
FOR USE ON SWEET CORN

ACCEPTED
FEB 14 1979
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pest(s)
registered under
EPA Reg. No. 352-342

Du Pont "Lannate" Insecticide is recommended for use on sweet corn for control of corn earworm. Control of earworm affecting ears is obtained through kill of eggs (ovicide) and larvae (larvicide).

HOW TO USE

Apply 1/4 to 1/2 lb. "Lannate" per acre using sufficient water to obtain uniform coverage (min. 3 gals. per acre by air). Apply when moths begin to lay eggs on silks and continue treatment at 1- to 3-day intervals.

Note: Do not graze or feed treated plants to livestock within 3 days after last application.

IMPORTANT

BEFORE USING "LANNATE", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

8/18/78

