


PLEASE NOTE

This image contains more than one label approved for this product on this date.

352-342

02/27/1979

1/1



AGRICULTURAL BULLETIN



LANNATE® INSECTICIDE
ALFALFA

ACCEPTED
FEB 27 1979
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under Reg. No. **352 342**

SUPPLEMENTAL LABELING
EPA REG. NO. 352-342

LANNATE® INSECTICIDE
FOR USE ON ALFALFA

Du Pont "Lannate" Insecticide is recommended for use on alfalfa for the control of loopers, yellowstriped armyworm and western yellowstriped armyworm.

HOW TO USE

Apply 1/4 to 1/2 lb. "Lannate" per acre when insects appear in damaging numbers; make repeat applications as needed. Use the higher rate for severe infestations.

Spray in sufficient water (min. 5 gals/per acre by air) to obtain thorough coverage.

- Note:
- 1) Do not apply to dormant or semi-dormant alfalfa when minimum daily temperature is 50° F. or lower.
 - 2) Do not graze or feed treated alfalfa to livestock within 7 days of last application.
 - 3) This product is toxic to bees and should not be applied when bees are actively visiting the area. In fields visited by honeybees, make application late in the evening or early morning. In areas frequented by alkali or leafcutting bees, do not apply during bloom or when the bees will be present in fields.

IMPORTANT

BEFORE USING "LANNATE", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

8/22/78

NEXT

LABEL

352-342

2/27/1979

AGRICULTURAL BULLETIN



LANNATE® METHOMYL INSECTICIDE
ANISE (FENNEL)



Under the Federal Insecticide, Fungicide, and Rodenticide Act, this product has been registered under the Federal Insecticide, Fungicide, and Rodenticide Act.
352-342
FEB 27 1979
ACCEPTED

SUPPLEMENTAL LABELING
EPA REG. NO. 352-342

LANNATE® METHOMYL INSECTICIDE
FOR USE ON ANISE (FENNEL)

Du Pont "Lannate" Methomyl Insecticide is recommended for use on anise (fennel) for control of cabbage looper and beet armyworm.

HOW TO USE

Apply "Lannate" at 5- to 7-day intervals or as needed in sufficient water (5 to 15 gal. per acre by air) to obtain uniform coverage.

- Beet armyworm - apply 1/2 to 1 lb. per acre
- Cabbage looper - apply 1 lb. per acre

Do not apply within 7 days of harvest.

Note: Pilot should not assist in the mixing and loading operation.

IMPORTANT

BEFORE USING "LANNATE", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental directions for use of this product which may not appear on package label. Follow the instructions carefully.

1/5/76