

PLEASE NOTE


This image contains more than one label approved for this product on this date.

352-342

7-26-1979



LANNATE® INSECTICIDE
COTTON



AGRICULTURAL BULLETIN

JUL
Under the Federal Insecticide, Fungicide, and Plant Disease Act, this product is registered under EPA Reg. No. 352-342

SUPPLEMENTAL LABELING
EPA REG. NO. 352-342

LANNATE® INSECTICIDE
FOR USE ON COTTON
WEST OF THE ROCKY MOUNTAINS

Du Pont "Lannate" insecticide is recommended for use on cotton to control tobacco budworm and cotton bollworm west of the Rocky Mountains. Control is obtained through kill of eggs (ovicide) and larvae (larvicide).

HOW TO USE

Apply 1/2 - 3/4 lb. "Lannate" per acre in sufficient water (use a minimum of 3 gals. per acre by air) to obtain thorough coverage. Repeat at 5- to 7-day intervals or as needed.

"Lannate" may be applied up to 15 days before harvest. Do not graze or feed treated cotton to livestock.

Note: Use of "Lannate" may redden cotton foliage. If reddening is excessive, alternate with other insecticides or discontinue use of "Lannate".

IMPORTANT

BEFORE USING "LANNATE", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

4/20/79



NEXT


LABEL

352-342

7-26-1979

LANNATE® INSECTICIDE
COTTON

AGRICULTURAL BULLETIN



SUPPLEMENTAL LABELING
EPA REG. NO. 352-342

LANNATE® INSECTICIDE
FOR USE ON COTTON
WEST OF THE ROCKY MOUNTAINS

Du Pont "Lannate" insecticide is recommended for use on cotton to control tobacco budworm and cotton bollworm west of the Rocky Mountains. Control is obtained through kill of eggs (ovicide) and larvae (larvicide).

HOW TO USE

Apply 1/2 - 3/4 lb. "Lannate" per acre in sufficient water (use a minimum of 3 gals. per acre by air) to obtain thorough coverage. Repeat at 5- to 7-day intervals or as needed.

"Lannate" may be applied up to 15 days before harvest. Do not graze or feed treated cotton to livestock.

Note: Use of "Lannate" may redden cotton foliage. If reddening is excessive; alternate with other insecticides or discontinue use of "Lannate".

IMPORTANT

BEFORE USING "LANNATE", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

4/20/79