

PLEASE NOTE

This image contains more than one label approved for this product on this date.

352-342

7-27-1979



AGRICULTURAL BULLETIN



LANNATE® INSECTICIDE
FALL ARMYWORM ON CROPS

SUPPLEMENTAL LABELING
EPA REG. NO. 352-342

LANNATE® INSECTICIDE
FOR CONTROL OF FALL ARMYWORM
ON CROPS

Du Pont "Lannate" Insecticide is recommended for use on cabbage, cucumber, potato, tomato, and tobacco to control fall armyworm.

HOW TO USE

Refer to the package label for spray volume and general use instructions.

Use Rates and Preharvest Limitations:

Crops	Lbs. "Lannate" Per Acre	Last Application - Days	
		To Harvest	To Livestock Grazing/Feeding
Cabbage	1/2	1	
Cucumber	1/2	1/2 lb. - 1; over 1/2 lb. - 3	
Potato	1/2	6	
Tomato	1/2	1/2 lb. - 1; over 1/2 lb. - 2	
Tobacco (Except shade)	1/2	5 flue cured 14 air or fire cured	

ACCEPTED
JUL 27 1979
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 352-342

IMPORTANT

BEFORE USING "LANNATE", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

5/1/79

NEXT

LABEL

352-342

7-27-1979



1/1



AGRICULTURAL BULLETIN

LANNATE® INSECTICIDE
COTTON
IN THE SOUTHWEST

SUPPLEMENTAL LABELING
EPA REG. NO. 352-342

LANNATE® INSECTICIDE
FOR USE ON COTTON
IN THE SOUTHWEST

Du Pont Lannate® insecticide is recommended for control of cotton fleahopper on cotton in the Southwest. The low use rates help conserve beneficial predators.

HOW TO USE

Apply 1/8 to 1/4 lb. of Lannate® per acre in sufficient water (minimum 2 gals. per acre by air) to obtain thorough coverage. Initiate schedule of sprays when significant numbers of fleahoppers are present. Continue at 5- to 7-day intervals.

Scouting of fields is required to insure proper timing of spray.

Note: Lannate® may be applied up to 15 days before harvest. Do not graze or feed treated cotton to livestock. Use of Lannate® may redden cotton foliage. If reddening is excessive, alternate with other insecticides or discontinue use.

IMPORTANT

BEFORE USING LANNATE®, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

ACCEPTED
JUL 27 1979
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
I am authorized to certify this pesticide
label as true and correct.
EPA Reg. No. 352-342

6/15/79


NEXT

LABEL

352-342

7-27-1979

1/1



AGRICULTURAL BULLETIN



LANNATE® INSECTICIDE
PEANUTS

SUPPLEMENTAL LABELING
EPA REG. NO. 352-342

LANNATE® INSECTICIDE
FOR USE ON PEANUTS

Du Pont "Lannate" Insecticide is recommended for use on peanuts for control of thrips.

HOW TO USE

Apply 1/2 to 1 lb. "Lannate" per acre as a spray in sufficient water to obtain thorough coverage (use a minimum of 3 gals. per acre for aerial application). Make up to 3 applications as needed to maintain control.

Note: Do not apply within 21 days of harvest.
Do not feed-treated vines to livestock.

IMPORTANT

BEFORE USING "LANNATE", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

5/22/79

ACCEPTED
JUL 27 1979
Under the Federal Insecticide
Fungicide and Rodenticide Act
as amended, this product is
registered under 352-342



NEXT

LABEL



352-342

7-27-1979

AGRICULTURAL BULLETIN



LANNATE® INSECTICIDE
PEANUTS

11

SUPPLEMENTAL LABELING
EPA REG. NO. 352-342

LANNATE® INSECTICIDE
FOR USE ON PEANUTS

Du Pont "Lannate" Insecticide is recommended for use on peanuts for control of granulate cutworm.

HOW TO USE

Apply 1/2 to 1 lb. "Lannate" per acre as a spray in sufficient water to obtain thorough coverage (use a minimum of 2 gals. per acre for aerial application).

For best results, treatment should be applied when cutworms are in the foliage of the crop; this occurs at night or on cloudy days.

Use up to three applications as needed to maintain control.

Note: Do not apply within 21 days of harvest.
Do not feed treated vines to livestock.

IMPORTANT

BEFORE USING "LANNATE", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

5/18/79

