




TERSAN[®]



WITH A ZINC SALT ADDED

ACTIVE INGREDIENT:

Maneb (Manganese ethylenebisdithiocarbamate) 80%
(Manganese equivalent as metallic) 16.5%)

INERT INGREDIENTS 20%

USDA Reg. No. 352-343

U.S. Pats. 2,710,822-3,379,610

Keep out of reach of children.

**CAUTION! MAY IRRITATE EYES, NOSE, THROAT, AND SKIN.
MAY BE HARMFUL IF INHALED OR SWALLOWED.**

Avoid breathing dust or spray mist. Keep away from fire or sparks.
Avoid contact with skin, eyes, and clothing.

In case of contact, flush skin or eyes with plenty of water; for eyes, get medical attention

IMPORTANT—Never allow "Tersan" LSR to become wet during storage. This may lead to certain chemical changes which will reduce the effectiveness of "Tersan" LSR. LSR is a fungicide and creates vapors which may be flammable. Keep container closed when not in use.

NET 3 LBS.

GENERAL INFORMATION—Du Pont "Tersan" LSR Turf Fungicide is recommended for use as a spray for the control of leaf spot diseases incited by *Helminthosporium* spp., rust, and *Rhizoctonia* brown patch on turfgrasses.

DIRECTIONS

"Tersan" LSR should be used only in accordance with recommendations on this label or in separate published Du Pont recommendations available through local dealers.

TURF DISEASES

Leaf Spot (*Helminthosporium* spp.)—Apply "Tersan" LSR at the rate of 3 to 4 oz. per 1000 sq. ft. using 3 to 5 gals. of water. Begin application when disease first appears and repeat at 7 to 10 day intervals as long as disease persists. When conditions are unusually favorable for disease activity, increase the rate to 6 to 8 oz. per 1000 sq. ft. and reduce interval to 3 to 5 days.

Rust—Apply "Tersan" LSR at the rate of 3 to 4 oz. per 1000 sq. ft. using 3 to 5 gals. of water when disease first appears and repeat at 7 to 10 day intervals as long as disease is evident.

Rhizoctonia brown patch—Apply "Tersan" LSR at the rate of 3 to 4 oz. per 1000 sq. ft. using 3 to 5 gals. of water when disease first appears and repeat at 7 to 10 day intervals as long as needed. When conditions are unusually favorable for disease activity, increase the rate to 6 to 8 oz. per 1000 sq. ft. and reduce interval to 3 to 5 days.

Do not re-use container, bury when empty. This product is toxic to fish. Do not dispose of any of this product or container in or near any body of water. Do not graze or feed clippings from treated areas to livestock.

NOTICE TO BUYER—Seller warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. This warranty does not extend to use of this product contrary to label use directions, or under abnormal use conditions, or under conditions not reasonably foreseeable to seller; buyer assumes all risk of any such use. Seller makes no other warranties, express or implied.

9-69

12-70

