

352-343 Pg. 1 of 2
PM 21

SPECIMEN LABEL
ACTUAL SIZE

ACCEPTED
MAY 15 1985
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 352-343



TERSAN®

Leaf Spot
Rust
Brown Patch

LSR

TURF FUNGICIDE

ACTIVE INGREDIENT

A coordination product of zinc ion and manganese
ethylene bisdithiocarbamate 80%
in which the ingredients are
Manganese⁺⁺ 16%
Zinc⁺⁺ 2%
Ethylene bisdithiocarbamate ion (C₂H₂N₂S₂)= 62%

INERT INGREDIENTS 20%

U.S. Pat. 3,379,610

EPA Est. 34302-CL-1

EPA Reg. No. 352-343

Keep out of reach of children.
PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
CAUTION! MAY IRRITATE EYES, NOSE, THROAT, AND SKIN.
MAY BE HARMFUL IF INHALED OR SWALLOWED.
Avoid breathing dust or spray mist. Avoid contact with skin, eyes, and clothing.
Keep away from fire or sparks.
Wear protective clothing including long pants, long sleeve shirt, gloves, hat and boots during mixing and loading operations.
First Aid: In case of contact, flush skin or eyes with plenty of water; for eyes, get medical attention.
ENVIRONMENTAL HAZARDS
This product is toxic to fish. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated.

NET 3 LBS.

MADE BY DU PONT IN COLOMBIA
E. I. DU PONT DE NEMOURS & CO. (INC.),
AGRICULTURAL CHEMICALS DEPARTMENT, WILMINGTON, DELAWARE

SPECIMEN LABEL

ACTUAL SIZE

Pg 2 of 2

NOTICE OF WARRANTY

Du Pont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Du Pont. In no case shall Du Pont be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. DU PONT MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

GENERAL INFORMATION—Du Pont "Tersan" LSR Turf Fungicide is recommended for use as a spray for the control of leaf spot diseases incited by *Helminthosporium* spp., rust, and *Rhizoctonia* brown patch on turfgrasses.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

"Tersan" LSR should be used only in accordance with recommendations on this label, or in separate published Du Pont recommendations available through local dealers.

Du Pont will not be responsible for losses or damages resulting from use of this product in any manner not specifically recommended by Du Pont. User assumes all risks associated with such nonrecommended use.

TURF DISEASES

Leaf Spot (*Helminthosporium* spp.), *Rhizoctonia* brown patch—Apply "Tersan" LSR at the rate of 3 to 4 oz. per 1000 sq. ft. using 3 to 5 gals. of water when disease first appears and repeat at 7 to 10 day intervals as long as needed. When conditions are unusually favorable for disease activity, increase the rate to 6 to 8 oz. per 1000 sq. ft. and reduce interval to 3 to 5 days.

Rust—Apply "Tersan" LSR at the rate of 3 to 4 oz. per 1000 sq. ft. using 3 to 5 gals. of water when disease first appears and repeat at 7 to 10 day intervals as long as disease is evident.

"Tersan" LSR plus Du Pont TERSAN® 1991 Turf Fungicide:

***Helminthosporium* leafspot plus dollar spot and/or brown patch; dollar spot plus brown patch; brown patch alone**—Apply 1 oz. "Tersan" 1991 plus 3-4 oz. "Tersan" LSR in 3 to 5 gallons of water for each 1,000 sq. ft. of turfgrass area to be treated. Make initial application when disease first appears and continue applications on 7- to 14-day intervals for as long as needed. Use the shorter intervals and higher use rate when conditions are unusually favorable for disease development. The combination of "Tersan" 1991 and "Tersan" LSR should not be used in areas where resistance to "Tersan" 1991 has occurred previously.

NOTE: Do not graze or feed clippings from treated areas to livestock.

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.

STORAGE AND DISPOSAL

STORAGE: Important—Never allow "Tersan" LSR to become wet during storage. This may lead to certain chemical changes which will reduce the effectiveness of "Tersan" LSR as a fungicide and create vapors which may be flammable. Keep container closed when not in use. Store product in original container only, away from other pesticides, fertilizer, food or feed.

DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.