

Du Pont will not be responsible for losses or damages resulting from use of this product in any manner not specifically recommended by Du Pont. User assumes all risks associated with such nonrecommended use.

Apply Hyvar XL before or during the period of active growth of plants to be controlled when rainfall can be expected for soil activation. If dense growth is present, results will be improved if vegetation is removed before treatment. Do not apply when ground is frozen.

Before spraying,ibrate equipment to determine quantity of water necessary to uniformly cover measured area to be treated. Measure the proper amount of Hyvar XL and mix into the necessary volume of water. After Hyvar XL has been thoroughly mixed in the spray tank, agitation of the spray solution is not required.

Use a hand gun power sprayer properly calibrated for constant speed and rate of delivery. For application with a hand gun sprayer, use at least 200 gals. of water per acre. For treating small areas, a tank type hand sprayer or sprinkling machine may be used. For basals, treatment of brush with Hyvar XL may be applied and diluted with an exact delivery hand gun applicator.

WEED CONTROL To control most weeds for an entire growing season at the end of non-cropland areas such as RAILROAD, HIGHWAY and POWER LINE RIGHT OF WAY, PETROLEUM TANK FARMS, LUMBER YARDS, STORAGE AREAS, and INDUSTRIAL PLANT SITES.

Apply 1 to 3 gals. Hyvar XL per acre for ANNUAL WEEDS and GRASSES such as foxtail, pigweed, wild oats, cat grass, nutgrass, bromegrass, ragweed, wild radish, purslane, chickweed, hardgrass and turkey mulch. When applied just prior to or after emergence of annuals, rates as low as 1 gal. per acre will control many annual weeds. Treat areas of low rainfall areas and areas of term control in higher rainfall areas.

Apply 3 to 5 gals. Hyvar XL per acre for PERENNIAL WEEDS and GRASSES such as bermudagrass, crabgrass, dogfennel, purple top, quackgrass, timothy, water hyacinth, and dogfennel. In high rainfall areas and wide brush areas with low rainfall, rates of 2 to 5 gals. per acre control many perennial weeds and grasses.

Apply 5 to 10 gals. Hyvar XL per acre for WOOLLY WINDMILL GRASS, use in the same manner as for PERENNIAL WEEDS and GRASSES such as bermudagrass, crabgrass, nutgrass, water hyacinth, dogfennel, quackgrass, timothy, and dogfennel. When applied in areas with less than 4 inches of rainfall, the effect will be improved in the areas of the West. Hyvar XL will control dogfennel, timothy, and quackgrass in the areas of the West.

Use the higher end of the dosage ranges in arid or semi-arid areas where high rainfall is not expected.

Retreatment Apply 1 to 3 gals. Hyvar XL per acre when annual weeds and grasses reappear in areas where weed growth has been controlled.

CONTINUED ON SIDE PANEL ▶

For Small Areas—Three fl. oz. of Hyvar XL per 250 sq. ft. is approximately 4 gals. per acre.

BRUSH CONTROL: To control undesirable woody plants on non-cropland areas such as RAILROAD RIGHT OF WAYS, STORAGE AREAS, INDUSTRIAL PLANT SITES and DRAINAGE DITCHES.

Apply as a broadcast or basal (soil) treatment for use on drainage ditches, apply as a basal (soil) treatment only. **Note:** For effective brush control and prevention of damage to desirable vegetation, do not apply to brush standing in water, do not use water from treated ditches for irrigation, do not use irrigation ditches for right of ways or other sites where marketable timber or other desirable trees or shrubs are immediately adjacent to the treated areas.

Broadcast Treatment Apply 2 to 3 gals. per acre to control oak, willow, sweet gum and pine at 1 to 1.2 gals. per acre to control brush such as American elm, winged elm, hackberry, sumac, and cottonwood. Use the higher levels of the dosage ranges on adsorptive soils (those high in organic matter or carbon).

Basal (Soil) Treatment Controls woody plants such as cottonwood, hackberry, maple, oak, poplar, red bud, sweet gum, wild cherry, willow, winged elm, American elm and sumac.

Diluted—Mix 1 gal. Hyvar XL in 5 gals. of water and apply at the rate of 1 to 2 fl. oz. per stem 2 to 4 in. basal diameter, wet base of stem to run off.

Undiluted—Apply Hyvar XL undiluted with an exact delivery hand gun applicator, this equipment delivers a thin stream of a predetermined volume when triggered. Apply at the rate of 5 to 10 ml. liters per stem 2 to 4 in. basal diameter, direct the treatment to a spot at the base of the trunk or collar area. When treating large stems and more than one delivery of solution is needed per stem, apply on the opposite side of the stem. **Note:** Hyvar XL is a combustible product, do not smoke while applying the undiluted product and keep away from heat and open flame.

NOTICE TO BUYER Purchase of this material does not confer any rights under patents in countries outside of the United States.

STORAGE AND DISPOSAL

STORAGE Store product in original container only away from other pesticides, fertilizer, food or feed. Keep container closed when not in use.

DISPOSAL Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at approved waste disposal facility. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

BEST AVAILABLE COPY