



# KROVAR<sup>®</sup> III

WEED KILLER

## WETTABLE POWDER

ACTIVE INGREDIENTS	80%
Bromacil [5-bromo-3-sec-butyl-6-methyluracil]	53%
Diuron [3-(3,4-dichlorophenyl)-1,1-dimethylurea]	27%
INERT INGREDIENTS	20%

Keep out of reach of children.  
**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS**  
**CAUTION!** MAY IRRITATE EYES, NOSE, THROAT, AND SKIN.  
 Avoid breathing dust or spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.  
**STATEMENT OF PRACTICAL TREATMENT**  
 Thoroughly clean all traces of "Krovar" II from application equipment immediately after use. Flush tank, pump, hoses, and boom with several changes of water after removing nozzle tips and screens (clean these parts separately).  
**ENVIRONMENTAL HAZARDS**

**IMPORTANT**—Injury to or loss of desirable trees or other plants may result from failure to observe the following

Do not apply (except as recommended for crop use), or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, tennis courts or similar areas. Do not use in home fruit plantings nor in citrus orchards interplanted to other trees or desirable plants. Prevent drift of dry powder or spray to desirable plants. Do not contaminate any body of water. Keep from contact with fertilizers, insecticides, fungicides and seeds.

Thoroughly clean all traces of "Krovar" II from application equipment immediately after use. Flush tank, pump, hoses, and boom with several changes of water after removing nozzle tips and screens (clean these parts separately).

### NET 6 LBS.

**E. I. du Pont de Nemours & Co. (Inc.)**  
**Biochemicals Department, Wilmington, Delaware**

### GENERAL INFORMATION

Du Pont Power II Herbicide is a wettable powder to be mixed in water and applied as a spray to control weeds in the yard and home crop areas.

Power II Herbicide is a selective herbicide which kills broad-leaved weeds and grasses. It is not effective against coniferous trees, shrubs, and other woody plants. It is not effective against annual weeds and grasses. It is not effective against perennial weeds and grasses. It is not effective against weeds and grasses which have been established in the yard and home crop areas.

Moisture is necessary to activate the herbicide. Best results are obtained if treatment is made to moist soil and moisture is supplied by rainfall or sprinkler irrigation within two weeks after application. The degree of control and duration of effect will vary with the amount applied, soil texture, rainfall, and other conditions.

### NOTICE OF WARRANTY

Du Pont warrants that this product conforms to the chemical description on the label thereon and is reasonably fit for the purpose stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Du Pont. In no case shall Du Pont be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. DU PONT MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

**NOTICE TO BUYER:** Purchase of this material is subject to the rights under patents of others outside of the United States.

### STORAGE AND DISPOSAL

- Prohibitions:** Do not use this material in or near water bodies, streams, rivers, or other bodies of water.
- Pesticide Disposal:** Do not pour or wash this material down the drain, into the street, or into any other drainage system.
- Container Disposal:** Do not reuse this material for any other purpose. Do not burn or incinerate this material. Do not use this material for any other purpose.

### DIRECTIONS FOR USE

For use in the yard and home crop areas, apply 1/2 to 1 lb. per acre. For use in the home crop areas, apply 1/2 to 1 lb. per acre. For use in the yard and home crop areas, apply 1/2 to 1 lb. per acre. For use in the home crop areas, apply 1/2 to 1 lb. per acre.

### CITRUS

Use only where trees have been established in the orchard for at least 1 year. Apply with a properly calibrated hand-pumped sprayer as a fan or broad cast treatment beneath and around the trees. Use sufficient water (10-40 gals. per acre) to obtain uniform coverage and contact of foliage and fruit with spray mist. Avoid over-spraying and shut off spray boom while starting, turning, slowing or stopping. Injury to trees may result from over-spraying in the spray boom. Do not spray during the rain or immediately after rain. Do not spray during the hot sun or during the heat of the day. Do not spray during the wind or during the rain.

**Perennial Weeds:** For perennial weeds, apply 1/2 to 1 lb. per acre. For annual weeds, apply 1/2 to 1 lb. per acre. For biennial weeds, apply 1/2 to 1 lb. per acre. For woody weeds, apply 1/2 to 1 lb. per acre.

Soil Texture	Lbs. Per Acre
Sand or sandy sand	4 to 6
Sandy loam	5 to 7
Silt loam, clay loam	6 to 8

Alternatively, make two applications of 1/2 to 1 lbs. per acre per year, apply in spring and summer. Partial control of perennials usually occurs with a single treatment; repeat applications are required to control perennials. Control may be improved by cultivation prior to treatment; otherwise, avoid working the soil as long as weed control continues. If weed effectiveness of the treatment may be reduced.

**Annual Weeds:** Apply 1/2 to 1 lb. per acre. For biennial weeds, apply 1/2 to 1 lb. per acre. For woody weeds, apply 1/2 to 1 lb. per acre. For perennial weeds, apply 1/2 to 1 lb. per acre.

**Seeding Vines (balsam apple and milkweed)—Florida:** Remove established vines from trees prior to treatment. Apply 1/2 to 1 lb. per acre with beginning of the spring or after cultivation in the spring, but before a new vine has formed. For best results, the treated area should extend at least 1 foot beyond the drip line of the tree. Additional applications should be made at 4 to 6 week intervals.

Apply 1/2 to 1 lb. per acre to control weeds in the yard and home crop areas. For use in the home crop areas, apply 1/2 to 1 lb. per acre. For use in the yard and home crop areas, apply 1/2 to 1 lb. per acre.

**Note:** Temporary injury of citrus leaves may occur following treatment. Because injury to citrus trees may result, do not use on soils that contain more than 1% of iron, aluminum, or calcium salts, nor on soils that are heavily covered or exposed to frost. Do not use on soils that are heavily covered or exposed to frost. Do not use on soils that are heavily covered or exposed to frost.



Do not re-plant treated areas in any crop within two years after last application. Injury to subsequent crops may result except that citrus trees may be planted one year after the last application.

### NON-CROP WEED CONTROL

For use in the yard and home crop areas, apply 1/2 to 1 lb. per acre. For use in the home crop areas, apply 1/2 to 1 lb. per acre. For use in the yard and home crop areas, apply 1/2 to 1 lb. per acre. For use in the home crop areas, apply 1/2 to 1 lb. per acre.

**Retreatment:** Apply 1/2 to 1 lb. per acre when annual weeds and grasses reappear in areas where weed control has been obtained.

**Small areas:** For use in the yard and home crop areas, apply 1/2 to 1 lb. per acre. For use in the home crop areas, apply 1/2 to 1 lb. per acre. For use in the yard and home crop areas, apply 1/2 to 1 lb. per acre. For use in the home crop areas, apply 1/2 to 1 lb. per acre.



75% OF ACTUAL SIZE