



AGRICULTURAL BULLETIN



BENLATE BENOMYL FUNGICIDE
MANGOES
8/8/72

SUPPLEMENTAL LABELING
EPA REG. NO. 352-354

BENLATE® BENOMYL FUNGICIDE
FOR CONTROL OF
CERTAIN DISEASES OF MANGOES

ACCEPTED

MAR 9 1973

UNDER THE
FEDERAL
FOR REGISTRATION
ED UNDER NO. 352-354

Du Pont "Benlate" Benomyl Fungicide is recommended for use on mangoes for control of anthracnose.

HOW TO USE

Apply 1 to 2 lbs. "Benlate" per acre with ground equipment in sufficient water to give thorough coverage of trees and fruit.

Begin applications at first appearance of panicles (approximately 2 inches long) and repeat at weekly intervals until all fruits are set. Continue to apply at 3 to 4 week intervals. Under conditions of severe disease pressure, use higher rate and closer interval. Do not apply within 14 days of harvest.

Note: Do not use alkaline pesticides, such as basic copper sulfate, Bordeaux mixture, or lime sulfur, as a tank mixture with "Benlate".

IMPORTANT

BEFORE USING "BENLATE", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental directions for use of this product which may not appear on package label. Follow the instructions carefully.

8/8/72