

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

JUL 27 1992

Ms. Janice K. Sharp, Ph.D.
E.I. Du Pont de Nemours & Co.
P. O. Box 30038
Wilmington, DE 19880-0038

Dear Ms. Sharp:

Subject: Proposed Label Amendment - Use Deletions
Du Pont Benlate Fungicide
EPA Registration No. 352-354
Your Submission Dated September 30, 1991

The label amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable provided you satisfactorily address and incorporate the following comments:

1. Under Environmental Hazards, revise the second sentence to read, "Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark, except for the registered aquatic use on rice."
2. The first paragraph under chemigation limits use to "crops specifically designated in directions for use on the particular crop." None of the directions for specific crops include recommendations for application by chemigation. Revise this paragraph to delete the restriction or indicate those crops intended to be treated by this method.
3. For peanuts, add "Do not make more than eight (8) applications per season."
4. The directions for use on rice must be revised. As written, only the precautionary statements appear to apply

CONCURRENCES							
SYMBOL							
SURNAME							
DATE							


-2-

to California. The limitation against use in California would be clearer if you add, "Do not apply to rice in California".

Enclosed is a stamped label for your records. Submit five (5) copies of the final printed label.

If you have any questions regarding this matter, please contact me or Sidney Jackson at (703) 305-7610.

Sincerely yours,


Susan Lewis
Product Manager (21)
Fungicide-Herbicide Branch
Registration Division (H7505C)

BEST AVAILABLE COPY

PH 21 352-354

REGISTRATION CODE 0391-002

PACKAGE SIZE
6 - 2 POUND BAGS / CARTON
6 - 5 POUND BAGS / CARTON



BENLATE®

FUNGICIDE
WETTABLE POWDER

ACTIVE INGREDIENT

Benomyl [Methyl 1-(butylcarbamoyl)-2-benzimidazolecarbamate]

BY WEIGHT

50%

INERT INGREDIENTS

50%

TOTAL

100% TLD

EXPIRATION DATE:
See Data Sheet

EPA Reg. No. 352-354

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION! MAY IRRITATE EYES, NOSE, THROAT AND SKIN.

Avoid breathing dust or spray mist. Avoid contact with skin, eyes, and clothing. Wash thoroughly after using.

Do not apply this product in such a manner as to directly or through drift expose workers, or other persons. The area being treated must be vacated by unprotected persons.

Protective clothing must be worn during periods of exposure. See Directions for Use.

This product may cause a temporary allergic skin reaction in a few susceptible persons. This condition should be treated as an allergic dermatitis. There is no evidence of aftereffects or permanent injury.

First Aid: In case of contact, flush skin or eyes with plenty of water; for eyes, get medical attention.

Aerial Mixer-Loaders: Wear a dust mask or a respirator during handling and mixing as this product may be harmful if inhaled. See Directions for Use.

For medical emergencies involving this product, call toll free 1-800-441-3537.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water. Do not apply where runoff is likely to occur. Drift and runoff from treated areas may be hazardous to fish in adjacent areas. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from areas treated.

For registered aquatic uses: Aquatic organisms may be killed at recommended application rates.

PHYSICAL OR CHEMICAL HAZARDS

Keep away from fire or sparks.

BEST AVAILABLE COPY

NOTICE

Do not enter or allow entry into treated areas for 24 hours after application unless wearing the protective clothing specified for this product or unless entering to perform post application activities on strawberries. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Warnings must include the following information. "CAUTION. Area treated with 'Benlate' on (date of application). Do not enter without appropriate protective clothing until sprays have dried. In case of contact, flush skin or eyes with plenty of water; for eyes, get medical attention."

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Du Pont "Benlate" should be used only in accordance with recommendations on this label, or in separate published Du Pont recommendations available through local dealers.

Du Pont will not be responsible for losses or damages resulting from use of this product in any manner not specifically recommended by Du Pont. User assumes all risk associated with such nonrecommended use.

"Benlate" is a systemic fungicide recommended for the control of many important plant diseases. If treatment is not effective following use of "Benlate" as recommended, a resistant strain of the fungus may be present. If treatment is ineffective due to the presence of a benomyl resistant strain, then neither "Benlate", nor any other benzimidazole or thiophanate type fungicide will effectively control that disease; consideration should be given to prompt use of other types of suitable fungicides.

The repeated exclusive use of "Benlate" may lead to buildup of resistant strains of fungi and loss of disease control. A spray program using other fungicides may delay resistant strain buildup. Consult your state extension specialist or official state recommendations for guidance on your particular crop and disease control situation.

NOTE: Do not tank mix or alternate "Benlate" with benzimidazole or thiophanate products such as "Merfex"[1] or "Topsin"[2].

Do not use on Greenhouse crops, including Hydroponic grown crops. Do not use on any container grown crops.

Do not enter or allow entry into treated areas for 24 hours after application unless wearing the protective clothing specified for this product or unless entering to perform post application activities on strawberries.

WORK SAFETY RULES

Keep all unprotected persons, children, livestock and pets away from treated area or where there is danger of drift.

During mixing and loading of the concentrated product wear the following protective clothing and equipment:

Long sleeve shirt and long legged pants; chemical resistant gloves, chemical resistant apron; chemical resistant shoes, shoe coverings, or boots. Wear a dust mask or a respirator approved by the National Institutes of Occupational Safety and Health (30 CFR Part 11) when mixing or loading for aerial application.

Wear the following protective clothing during application, equipment repair, equipment cleaning, and during reentry to treated areas before expiration of the 24-hour reentry interval:

Long sleeve shirt and long legged pants; chemical resistant gloves; chemical resistant shoes, shoe coverings, or boots. During aerial application from enclosed cockpits wear long sleeve shirt and long legged pants. A clean pair of chemical resistant gloves must be carried in the cockpit and worn during exiting from the aircraft.

Important! Before removing gloves, wash them with soap and water. Always wash hands, face, and arms with soap and water before smoking, eating, drinking or toileting.

After work, take off all clothes and shoes. Shower using soap and water. Wear only clean clothes. Do not use contaminated clothing. Wash

protective clothing and protective equipment with soap/detergent and water after each use. Personal and protective clothing worn during use must be laundered separately from household articles. Clothing or protective equipment heavily contaminated or drenched with benomyl must be destroyed according to state or local regulations. Heavily contaminated or drenched clothing cannot be adequately decontaminated.

Flaggers must be in totally enclosed vehicles.

Apply as a spray with ground equipment (except as otherwise directed), using sufficient water to obtain thorough coverage of plants. Under severe disease conditions use the higher rate and shorter interval specified for each crop; also, for tree crops, use the higher rate for large mature trees. For aerial application (listed crops only) use the following gals. per acre: Rice and Soybeans, 3 to 10; Cabbage (seed crop), Celery, Cucurbits, Peanuts and Sugar Beets, 5 to 10; Almonds, Avocados, Beans, Pecans, Stone Fruits and Strawberries, 10 to 20; Grapes, 15 to 20.

For use in small gardens and orchards (less than 1 acre), application rates may be converted to lbs. per 100 gals. by dividing the lbs. per acre rate in half, and applying the resulting spray mixture at the rate of 4.5 gals. per 100 sq. ft. (1 lb. "Benlate" per 100 gals. equals 1 tablespoonful per gal.)

Add required amount of "Benlate" to necessary volume of water in spray tank agitated by hydraulic or mechanical means; continuous agitation is required to keep the material in suspension. Do not tank mix "Benlate" with lime or alkaline pesticides such as Bordeaux mixture or lime sulfur.

Where use of spray oil is recommended (apples, peanuts, pecans, stone fruits), use a nonphytotoxic superior-type (60 to 70 second viscosity) spray oil; add as last ingredient to spray tank. Before applying other pesticides in conjunction with spray oil or immediately before or after oil application, consult product labels. Observe all cautions and limitations on labeling of all products used in mixtures.

Grazing or feeding livestock is prohibited on forage and hay of benomyl treated wheat, and on forage and hay of barley, oats, rye and wheat following seed treatment.

CHEMIGATION: Apply "Benlate" only through sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set or hand move irrigation systems only on crops specifically designated in the "Directions for Use" for that particular crop. Do not apply "Benlate" through any other type of irrigation system or on any other crops.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service Specialists, equipment manufacturers or other experts.

Do not connect an irrigation system used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Specific Instructions for Public Water Systems:

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

Specific Instructions for Sprinkler Irrigation Systems:

1. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.
8. Good agitation is required in the injection tank.

9. In moving systems, apply specified dosage of "Benlate" as a continuous injection. In nonmoving systems inject "Benlate" for 15 to 30 minutes at end of cycle. Use the least amount of water possible consistent with uniform coverage.
10. Mix the amount of "Benlate" needed for acreage to be treated into the quantity of water determined during prior calibration. For moving systems inject into the system continuously for one complete revolution of the field. For nonmoving systems inject into system for the time established during calibration.
11. Stop injection equipment after treatment is completed and continue to operate irrigation equipment until all "Benlate" is flushed from system.

ALMONDS: Brown Rot Blossom Blight - Apply 1 to 1 1/2 lbs. per acre at pink bud. Under severe disease conditions and on highly susceptible varieties, make a second application during half-to-full bloom.

APPLES: For applications through cover sprays, use "Benlate" as a tank mixture as detailed below. Apply 200 to 500 gals. of spray per acre with hydraulic ground equipment or equivalent amount of products per acre with concentrate sprayers. Do not graze livestock in treated orchards.

"Benlate" + Captan: Scab, Powdery Mildew, Sooty Blotch, Flyspeck, Bitter Rot, Black Rot - Use 2 to 3 ozs. "Benlate" plus 12 to 16 ozs. Captan 50WP Fungicide (or 7 1/2 to 10 ozs. Captan 80WP) per 100 gals. of water. Apply at 1/2" green tip and repeat at 7- to 14-day intervals (or as needed) through the cover sprays. Use the 3 oz. rate of "Benlate" for varieties more susceptible to powdery mildew. If an application is missed during an infection period, apply the higher rates as soon as possible after the infection period in order to deactivate scab and to prevent further infection. NOTE: Spray injury may result if Captan is used with, immediately before, or closely following an oil spray.

Postharvest Fruit Rots (*Botrytis* spp., *Penicillium* spp. *Gloeosporium* spp.) - Make a single application of 6 ozs. "Benlate" per 100 gals. anytime from 3 weeks before harvest up to day of harvest.

AVOCADOS (Florida): Scab, *Cercospora* Spot, Anthracnose - Apply 1 to 2 lbs. per acre; begin when buds swell and repeat at 3- to 4-week intervals. Do not apply within 30 days of harvest.

BEANS: White Mold (*Sclerotinia*), Gray Mold, (*Botrytis*) - Use on beans grown as fresh vegetables, for processing, or for the dry bean market. Apply 1 1/2 to 2 lbs. per acre at 25% to 50% bloom; repeat at peak bloom. For narrow-row (20-40") irrigated dry beans in Montana, Nebraska, Colorado and Wyoming, apply at initial bloom and repeat 7 to 10 days later; only partial control of white mold may result.

Do not apply within 14 days of harvest (28 days for lima beans), do not use where crop is grown only for forage purposes.

BLUEBERRIES Apply 1 lb. per acre. Do not make more than 4 applications before harvest; do not apply within 21 days of harvest. Mummy Berry, Botrytis Blossom Blight - Apply at green tip and repeat at 7- to 10-day intervals through petal fall. Anthracnose Leafspot - Apply when disease first appears and make one additional application 14 days later. After harvest, make up to 4 applications to the bushes at 14-day intervals as needed. Do not use on container grown blueberries.

CABBAGE (Seed Crop - Pacific Northwest) White Blight (*Sclerotinia* Stalk Rot) - Apply 2 lbs. per acre by aircraft in 5 to 10 gals. of water; add a spreader-sticker to aid in wetting plants. Make first application at first petal fall; make two additional applications at 14-day intervals if conditions favor development of disease. Note: Do not graze treated areas; do not use seed or plant parts for food or feed purposes.

CANEBERRIES: RASPBERRIES, BLACKBERRIES, BOYSENBERRIES, LOGANBERRIES, DEWBERRIES: Botrytis, Powdery Mildew, Penicillium Rots - Apply 3/4 lb. per acre at early bloom (5 to 10%) and at full bloom; make up to 3 additional applications at 14-day intervals as needed. Do not apply within 3 days of harvest.

CELERY: Early Blight (*Cercospora*), Late Blight (*Septoria*) - Apply 1/4 to 1/2 lb. per acre; begin when disease first appears and repeat at 7- to 10-day intervals. Do not apply within 7 days of harvest.

CITRUS: Scab - Apply 1 1/2 to 3 lbs. per acre. Under conditions of severe disease pressure, apply at pinhead stage (just prior to first flush) and repeat at 2/3 petal fall; otherwise, make a single application at 2/3 petal fall.

Greasy Spot - Make a single application of 1 1/2 to 3 lbs. per acre during the period mid-June to mid-July.

Fruit Decay (Green Mold, Blue Mold, Stem-end Rot) - Make a single application of 1 to 2 lbs. per acre anytime from 3 weeks prior to harvest up to day of harvest.

Note: Do not graze livestock in treated groves.

CUCURBITS: CUCUMBERS, MELONS, PUMPKINS, SUMMER AND WINTER SQUASH: Target Spot (Cucumbers), Gummy Stem Blight, Powdery Mildew, Anthracnose - Apply 1/4 to 1/2 lb. per acre; for aerial application, use 1/2 lb. per acre. Begin applications when plants begin to run or when disease first appears, and repeat at 7- to 14-day intervals as needed. For target spot, use 7-day intervals as needed.

GRAPES: Botrytis Bunch Rot - Apply 1 to 1 1/2 lbs. per acre at first bloom (no later than 5% bloom) and repeat 14 days later if severe disease conditions persist. Make an additional application 3 to 4 weeks before harvest or when sugar begins to build; repeat 14 days later if conditions favorable for disease persist. "Benlate" does not control bunch rots caused by other organisms such as *Rhizopus* spp., *Alternaria* spp., and *Diplodia* spp.; these rots occur most frequently in high temperature areas such as the San Joaquin and Sacramento Valleys of California. Powdery Mildew, Black Rot, Bitter Rot - East of Rockies - Apply 3/4 to 1 1/2 lbs. per acre when foliage first develops and repeat at 14- to 21-day intervals, or as needed, until berries are full size.

Note: Do not apply within 7 days of harvest.

MACADAMIA NUTS (Hawaii): Botrytis Blossom Blight - Apply 1 3/4 lbs. per acre; a surfactant may be added to the spray to improve wetting of foliage. Begin applications 1 to 2 weeks prior to bloom, and repeat at 7- to 14-day intervals through the bloom period.

MANGOES: Anthracnose - Apply 1 to 2 lbs. per acre. Begin applications at first appearance of panicles (approx. 2" long), and repeat at weekly intervals until all fruits are set. Continue at 3- to 4-week intervals. Do not apply within 14 days of harvest.

MUSHROOMS: Verticillium Spot (Dry Bubble) - Use 1 lb. per 100 gals. and apply to bed surface at the rate of 12 1/2 gals. per 1000 sq. ft. Apply immediately after casing and repeat at pinning; alternatively, if disease has occurred, apply to beds after picking and repeat 10 days later. Do not apply within 2 days of harvest.

PEANUTS: "Benlate" + "Manzate" 200 Fungicide: *Cercospora* Leafspot, Rust, *Ascochyta* Web Blotch - Apply 1/4 lb. "Benlate" plus 1 1/2 lbs. "Manzate" 200 per acre; spray oil may be added at the rate of 1 pt. to 1 qt. per acre. Begin applications 35 to 40 days after planting or when disease first appears. Repeat at following intervals: for *Cercospora* leafspot, 10 to 14 days; for rust, 7 to 14 days; for *ascochyta* web blotch, 7 to 14 days. Do not apply within 14 days of harvest; do not graze or feed treated vines, hay, or hulls to livestock.

PEARS: Scab, Powdery Mildew, Sooty Blotch, Flyspeck - Use 4 to 6 ozs. per 100 gals. of water, apply 200 to 500 gals. of spray per acre with hydraulic ground equipment or equivalent amount of "Benlate" per acre with concentrate sprayers. Apply at 1/2" green tip and repeat at 7- to 14-day intervals (or as needed) through the cover sprays. If an application is missed during an infection period, use 6 ozs. per 100 gals. and apply as soon as possible after the infection period in order to deactivate scab and to prevent further infection. Do not graze livestock in treated orchards.

Postharvest Fruit Rots (Botrytis spp., Penicillium spp., Gloeosporium spp.) - Make a single application of 6 ozs. per 100 gals. anytime from 3 weeks before harvest up to day of harvest.

Overwintering Scab - Apply 8 ozs. per 100 gal. after harvest but before leaf drop. Thorough wetting of foliage is necessary.

PECANS: Pecan Scab, Brown Leafspot, Downy Spot, Powdery Mildew, Liverspot, Zonate Leafspot, Fungal Leaf Scorch - Apply 1/2 to 1 lb. per acre; use the higher rate on trees over 30 ft. tall. For aerial application (Ark., La., Miss., Okla., Tex., only), use 1 lb. per acre. Spray oil may be added at the rate of 1 to 2 gals. per acre. Apply at prepollination when young leaves are unfolding, when small nuts are forming, and thereafter at 3- to 4-week intervals. Do not apply after shucks split.

PINEAPPLE: Pineapple Butt Rot (*Thielaviopsis paradoxa*) - Use 1 1/4 lbs. per 100 gals. of water as a pre-plant dip treatment. Immerse seedpieces to give thorough wetting; remove and allow to drain.

RICE (Except California) - Do not apply within 21 days of harvest. Do not apply to stubble rice. Do not apply to fields where crayfish or catfish farming is practiced, nor drain water from treated areas into areas where such farming is practiced. Water drained from treated areas must not be used to irrigate other crops.

Rice Blast, Stem Rot - Apply 1 to 2 lbs. per acre at booting and repeat at heading.

SOYBEANS: Do not apply within 35 days of harvest; do not graze or feed treated soybean vines or hay to livestock.

Diaporthe Pod-and-Stem Blight, Anthracnose, Septoria Brown Spot, Cercospora Frogeye Leafspot, Purple Seed Stain - Apply 1/2 to 1 lb. per acre. For determinate varieties (generally grown in the South), apply at early pod set when majority of pods are 1/8 to 1/2" in length, for indeterminate varieties (generally grown in the North), apply when pods near the top of the plant are 1/2 to 1" in length. Make one additional application 14 to 21 days later.

STONE FRUITS: APRICOTS, CHERRIES, NECTARINES, PEACHES, PLUMS, PRUNES: Treatment is most effective if applied just before rainfall; for aerial application, fly over every row or center.

EAST OF ROCKY MOUNTAINS: Use 3/4 to 1 1/2 lbs. per acre on trees up to 12 ft. tall; over 12 ft., use 1 1/2 to 2 lbs. per acre.

Brown Rot Blossom Blight - Apply at early bloom stages (apricots - red bud; peaches, nectarines - pink bud; cherries - early popcorn, plums and prunes - green tip); for this application only, "Benlate" may be used in combination with spray oil. Make a second application at 75% to 100% bloom. If blossoming is prolonged or conditions favorable for disease persist, apply at petal fall.

Fruit Brown Rot - After blossom blight sprays, make two preharvest applications beginning 3 weeks before harvest up to day of harvest.

Peach Scab, Powdery Mildew - Use same schedule as for Brown Rot Blossom Blight plus applications at shuck split, shuck fall and 14 days later.

Cherry Leaf Spot - Use same schedule as for Brown Rot Blossom Blight and continue at 10- to 14-day intervals through harvest. Make an additional application 2 to 3 weeks after harvest.

WEST OF ROCKY MOUNTAINS - Use 1 1/2 to 2 lbs. per acre.

Brown Rot Blossom Blight - Apply at early bloom stages (apricots - red bud; peaches, nectarines - pink bud; cherries - early popcorn; plums and prunes - green tip); for this application only, "Benlate" may be used in combination with spray oil. If blossoming is prolonged or conditions favorable for disease persist, make a second application 14 days later.

Fruit Brown Rot - After blossom blight sprays, make a preharvest application (before rain) any time from 3 weeks before harvest to day of harvest. Make a second application if conditions favorable for disease persist or harvest is prolonged. Preharvest applications are most effective when applied with ground equipment, using sufficient volume to provide thorough and uniform coverage of fruit.

Powdery Mildew - Use same schedule as Brown Rot Blossom Blight plus applications at shuck split, shuck fall, and 14 days later.

Cherry Leafspot - Use same schedule as for Brown Rot Blossom Blight and continue at 10- to 14-day intervals through harvest. Make an additional application 2 to 3 weeks after harvest.

NOTE: "Benlate" does not control peach leaf curl, shot hole (*Coryneum* blight) or bacterial blast, nor fruit rots caused by *Rhizopus* spp. and *Alternaria* spp. Do not graze livestock in treated orchards.

STRAWBERRIES: Gray Mold (*Botrytis*) Powdery Mildew, Leaf Scorch, Leaf Blight, Leafspot - Apply 1 lb. per acre at 10% bloom and at full bloom; continue at 10- to 14-day intervals, using 1/2 lb. per acre. Anthracnose - Apply 1 lb. per acre when plants are established and repeat at 7-day intervals.

SUGARBEETS: *Cercospora* Leafspot - Apply 3/8 to 1/2 lb. per acre. Begin application when disease first appears and repeat at 14- to 21-day intervals as needed. Do not apply within 21 days of harvest.

TOMATOES - Field: Gray Mold (*Botrytis*), Leaf Mold (*Cladosporium*), White Mold (*Sclerotinia*), *Cercospora* Leafspot, *Phoma* Leafspot - Apply 1/2 to 1 lb. per acre. Begin applications when disease first appears and repeat at 7- to 14-day intervals as needed.

STORAGE AND DISPOSAL

STORAGE: Never allow "Benlate" to become wet during storage. This may lead to certain chemical changes which will reduce the effectiveness of "Benlate" as a fungicide. Keep container tightly closed when not in use. Store product in original container only, away from other pesticides, fertilizer, food or feed.

PRODUCT DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

NOTICE OF WARRANTY

Du Pont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Du Pont. In no case shall Du Pont be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. DU PONT MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

1 Registered trademark of Merck & Company.

2 Registered trademark of Nippon Soda Company, Japan

AG-5386 9031/9091/9101 3/27/91

BEST AVAILABLE COPY