



PM 21 352-354

SUPPLEMENTAL LABELING

10/1

DuPont Agricultural Products



BENLATE® FUNGICIDE
GRAPES
CA AND AZ

..... A Growing Partnership With Nature

BENLATE® FUNGICIDE

(EPA Reg. No. 352 - 354)

FOR TABLE GRAPES, WINE/JUICE GRAPES, AND GRAPES GROWN FOR RAISINS
IN THE STATES OF ARIZONA AND CALIFORNIA

DIRECTION FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Crop	Disease	Limit /Acre /Crop	Rate, Minimum Gallonage	Initial Application	Repeat Application	PHI
Grapes*	Botrytis Bunch Rot	96 oz (6 lb)	16 - 24 oz/A (1 - 1.5 lb/A) 15 gal/A air	First bloom (1-5%)	14 days after first bloom; Additional sprays 3-4 weeks before harvest (sugar increase) and 14 days later if conditions favor disease.	7 days
	Anthracnose (<i>Elsinoe</i>) Isariopsis Leaf Spot		24 oz/A (1.5 lb/A) 15 gal/A air			

- * "Benlate" does not control Rhizopus, Alternaria, or Diplodia Bunch Rots.
- * These rots occur most frequently in high temperature areas such as San Joaquin and Sacramento Valleys of California.
- * Do not use "Benlate" alone. Use only in combination with labeled non-Benzimidazole fungicide.

IMPORTANT


BEFORE USING "BENLATE", READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA REGISTERED LABEL.

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

011193
020894

04/07/93



BEST AVAILABLE COPY