

BENLATE FUNGICIDE

EPA REG. NO. 352-354

FOR THE SUPPRESSION OF ASHY STEM BLIGHT (Macrophomina phaseolina) OF SNAP BEANS IN THE STATE OF TEXAS

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

HOW TO USE

Apply 16 to 24 oz "Benlate" per acre, using sufficient water to obtain thorough coverage; a minimum of 15 gallons per acre for ground application, and 5 gallons per acre for aerial application. Apply post-emergence when conditions are right for infection. If needed, make a second application anytime prior to bloom.

Apply no more than a total of 4 lb "Benlate" per acre per crop.

Make no "Benlate" applications within 14 days of harvest.


IMPORTANT

BEFORE USING BENLATE, READ AND FOLLOW ALL APPLI-CABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

D-091796/4



RECD EPA/OPP/OPD 瓶 3 P2 ž

ģ

© 1996 E. I. du Pont de Nemours and Company, DuPont Agricultural Products, Wilmington, Delaware 19898

Page 1 of

Н-