

Ms. Janice K. Sharp, Ph.D.
E.I. Du Pont de Nemours & Co.
P. O. Box 80038
Wilmington, DE 19880-0038

Dear Ms. Sharp:

Subject: Proposed Label Amendment - Use Deletions
Du Pont Tersan 1991 Turf Fungicide
EPA Registration No. 352-357
Your Submission Dated September 30, 1991

The label amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable provided you satisfactorily address and incorporate the following comments:

1. Under Environmental Hazards, revise the second sentence to read, "Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark."
2. In the first sentence under chemigation, delete "only on crops specifically designated in directions for use for that particular crop." In the second sentence, delete "or on any other crop."

Enclosed is a copy of the stamped label. Submit five (5) copies of the final printed label.

If you have any questions regarding this matter, please contact me or Sidney Jackson at (703) 305-7610.

Sincerely yours,

SC

BEST AVAILABLE COPY

Susan Lewis
Product Manager (21)
Fungicide-Herbicide Branch
Registration Division (H7505C)

CONCURRENCES							
SYMBOL							
SURNAME							
DATE							



Tersan[®] 1991

TURF FUNGICIDE



TRADEMARK

WETTABLE POWDER



Tersan® 1991

TURF FUNGICIDE

ACTIVE INGREDIENT

Benomyl [Methyl 1-(butylcarbamoyl)-2-benzimidazolecarbamate] 50%

INERT INGREDIENTS

..... 50%

EPA Reg. No. 352-357

KEEP OUT OF REACH OF CHILDREN PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION! MAY IRRITATE EYES, NOSE, THROAT AND SKIN.

Avoid breathing dust or spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after using.

Do not apply this product in such a manner as to directly or through drift, expose workers or other persons. The area being treated must be vacated by unprotected persons.

Protective clothing must be worn during periods of exposure. See Directions for Use.

This product may cause a temporary allergic skin reaction in a few susceptible persons. This condition should be treated as an allergic dermatitis. There is no evidence of aftereffects or permanent injury.

First Aid: In case of contact, flush skin or eyes with plenty of water, for eyes, get medical attention.

For medical emergencies involving this product, call toll free 1-800-441-3637.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water. Do not apply where runoff is likely to occur. Drift and runoff from treated areas may be hazardous to fish in adjacent areas. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from areas treated.

PHYSICAL OR CHEMICAL HAZARDS

Keep away from fire or sparks

STORAGE AND DISPOSAL

STORAGE: Never allow "Tersan" 1991 to become wet during storage. This may lead to certain chemical changes which will reduce the effectiveness of "Tersan" 1991 as a fungicide. Keep container closed when not in use. Store product in original container only, away from other pesticides, fertilizer, food or feed.

DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

NOTICE OF WARRANTY

Du Pont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Du Pont. In no case shall Du Pont be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. DUPONT MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Du Pont Tersan 1991 should be used only in accordance with recommendations on this label, or in separate published Du Pont recommendations available through local dealers.

Du Pont will not be responsible for losses or damages resulting from use of this product in any manner not specifically recommended by Du Pont. User assumes all risks associated with such nonrecommended use.

Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Warnings must include the following information: "CAUTION: Area treated with Tersan 1991 on (date of application). Do not enter without appropriate protective clothing until sprays have dried. In case of contact, flush skin or eyes with plenty of water, for eyes, get medical attention."

"Tersan" 1991 Turf Fungicide is recommended for use as a spray for the control of dollar spot, large brown patch, Fusarium patch (pink snow mold), Fusarium blight and stripe smut on golf courses, parks, athletic fields, and commercial turf areas. ~~It is also recommended for the control of certain diseases of roses, flowers, ornamentals, shade trees, and bulbs.~~

If treatment is not effective following use of "Tersan" 1991 as recommended, a resistant strain of the fungus may be present. If treatment is ineffective due to the presence of a benomyl resistant strain, then neither "Tersan" 1991 nor any other benzimidazole or thiophanate type fungicide will effectively control that disease, consideration should be given to prompt use of other types of suitable fungicides.

The repeated exclusive use of "Tersan" 1991 may lead to buildup of resistant strains of fungi and loss of disease control. A spray program using other fungicides may delay resistant strain buildup. Consult your state extension specialist or official state recommendations for guidance on your particular crop and disease control situation.

NOTE: Do not tank mix or alternate "Tersan" 1991 with benzimidazole or thiophanate products such as "Mertect", "Topsin", "Fungo", or "3336".

WORK SAFETY RULES

Keep all unprotected persons, children, livestock and pets away from treated area or where there is danger of drift.

During mixing and loading of the concentrated product wear the following protective clothing and equipment:

Long sleeve shirt and long legged pants, chemical resistant gloves, chemical resistant apron, chemical resistant shoes, shoe coverings, or boots.

Wear the following protective clothing during application, equipment repair and equipment cleaning:

Long sleeve shirt and long legged pants, chemical resistant gloves, chemical resistant shoes, shoe coverings, or boots.

Important! Before removing gloves, wash them with soap and water. Always wash hands, face, and arms with soap and water before smoking, eating, drinking, or toileting.

After work, take off all clothes and shoes. Shower using soap and water. Wear only clean clothes. Do not use contaminated clothing. Wash protective clothing and protective equipment with soap, detergent and water after each use. Personal and protective clothing worn during use must be laundered separately from household articles. Clothing or protective equipment heavily contaminated or drenched with benomyl must be destroyed according to state or local regulations. Heavily contaminated or drenched clothing cannot be adequately decontaminated.

CHEMIGATION: Apply "Tersan" 1991 Turf Fungicide only through sprinkler including center pivot, lateral move, end tow, side (wheel on) traveler, big gun, solid set or hand move irrigation systems only on crops specifically designated in the "Directions For Use" for that particular crop. Do not apply "Tersan" 1991 through any other type of irrigation system or on any other crops.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service Specialists, equipment manufacturers or other experts.

Do not connect an irrigation system, including greenhouse systems, used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation systems and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Specific Instructions for Public Water Systems:

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

Specific Instructions for Sprinkler Irrigation Systems:

1. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

BEST AVAILABLE COPY

- 6 The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected
- 6 Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock
- 7 Do not apply when wind speed favors drift beyond the area intended for treatment
- 8 Good agitation is required in the injection tank
- 9 In moving systems, apply specified dosage of "Tersan" 1991 as a continuous injection. In nonmoving systems inject "Tersan" 1991 for 15 to 30 minutes at end of cycle. Use the least amount of water possible consistent with uniform coverage
- 10 Mix the amount of "Tersan" 1991 DF needed for acreage to be treated into the quantity of water determined during prior calibration. For moving systems inject into system continuously for one complete revolution of the field. For nonmoving systems inject into system for the time established during calibration
- 11 Stop injection equipment after treatment is completed and continue to operate irrigation equipment until all "Tersan" 1991 DF is flushed from system

TURF—NOTE The repeated and exclusive use of "Tersan" 1991 DF has led to the development of resistant strains of dollar spot in some locations and may lead to the development of resistance in other strains of fungi. The use of "Tersan" 1991 DF as a tank mixture with Daconil® 2787 Fungicide, Daconil® 2787 Flowable Fungicide, or Du Pont TERSAN® LSR Turf Fungicide may delay the development of resistant strains of fungi. Apply recommended amounts of "Tersan" 1991 DF plus either "Daconil" 2787, or "Tersan" LSR in sufficient water to obtain thorough coverage, usually 3 to 5 gals per 1,000 sq ft. When treating golf greens, always treat aprons and approaches.

"Tersan" 1991 plus "Daconil" 2787: Dollar Spot and Brown Patch—Apply 1 oz "Tersan" 1991 plus 3 fl oz "Daconil" 2787 Flowable (or 2 oz Daconil 2787 75% wettable powder) per 1,000 sq ft. Make the initial application when conditions favor disease development and repeat the treatment on 10- to 14-day intervals as needed.

"Tersan" 1991 plus "Daconil" 2787: Helminthosporium leafspot and Brown Patch (and Dollar Spot, if present)—Apply 1 oz "Tersan" 1991 plus 3 to 6 fl oz "Daconil" 2787 Flowable (or 2 to 4 oz "Daconil" 2787 75% wettable powder) per 1,000 sq ft. Make the initial application when conditions favor disease development and repeat the treatment on 7- to 10-day intervals as needed. Use the higher rate of "Daconil" 2787 in the combination if disease conditions are particularly severe.

"Tersan" 1991 plus "Tersan" LSR: Helminthosporium leafspot and dollar spot and/or brown patch; dollar spot and brown patch; brown patch—Apply 1 oz "Tersan" 1991 plus 3-4 ozs "Tersan" LSR for each 1,000 sq ft of turfgrass area to be treated. Make initial application when disease first appears and continue applications on 7- to 14-day intervals for as long as needed. Use the shorter intervals and higher use rate when conditions are unusually favorable for disease development.

"Tersan" 1991 in other spray programs—Apply recommended amounts of "Tersan" 1991 in sufficient water to obtain thorough coverage, usually 2 to 5 gals per 1,000 sq ft. When treating golf greens, always treat aprons and approaches.

Dollar Spot (*Sclerotinia homoeocarpa*)—Apply at rate of 1 oz per 1,000 sq ft at earliest appearance of disease and continue at 10- to 14-day intervals as long as needed.

Large Brown Patch (*Rhizoctonia solani*)—Apply at rate of 2 oz per 1,000 sq ft at earliest appearance of disease and continue at 10- to 14-day intervals as long as needed. When conditions are unusually favorable for development of disease, reduce interval to 5 to 7 days.

Fusarium Patch (*Fusarium nivale*)—A cool season disease. Apply at rate of 2 oz per 1,000 sq ft at earliest appearance of disease and continue at 10- to 14-day intervals as long as needed.

Fusarium Blight (*Fusarium roseum* and *F. tricinctum*)—A warm season disease. Apply at rate of 5 to 8 oz per 1,000 sq ft at earliest appearance of disease and repeat 10 to 14 days later. Immediately after application, water sufficiently to thoroughly wet soil to a depth of 1 inch below any mat or thatch present.

Stripe Smut (*Ustilago striiformis*)—Make a single application at rate of 6 oz per 1,000 sq ft in October or in early spring before grass begins growth. After application, water turf sufficiently to carry product to base of plants.

NOTE: Where Helminthosporium diseases are a problem, apply "Tersan" LSR Turf Fungicide two or three times in the early spring at rate of 3 to 4 oz per 1,000 sq ft at 7- to 10-day intervals.

Do not graze or feed clippings from treated areas to livestock

delete

ROSES, FLOWERS, ORNAMENTALS, SHADE TREES—Field and Greenhouse: Foliar Spray—Begin applications when disease first appears and repeat at 10- to 14-day intervals throughout the growing season, shorten interval during humid, rainy weather. Use at the following rates: **1 lb per 100 gals** (1 tablespoonful per 2 gals) — for Powdery Mildew, Botrytis Gray Mold, **1 lb per 100 gals** — for Anthracnose (for shade trees and woody ornamentals, begin at bud break and make 2 or 3 additional applications at 10- to 14-day intervals). Black Spot of roses, Cercospora, Entomosporium, Ramularia, and Septoria Leafspots, Ascochyta and Phomopsis blights, Didymella Leafspot of iris, Cornynespora Leafspot of Ligustrum, Oculmyces Blight of azalea and rhododendron (begin as flowers open). Scab of pyracantha and flowering crab. Addition of a surfactant to the spray mixture improves distribution of the spray on hard-to-wet plants such as roses. For aerial application, use 1 to 1.5 lb per acre.

Drench Treatment—Botrytis, Fusarium, Rhizoctonia and Sclerotinia stem, crown and root rots on herbaceous annuals, perennials and bedding plants. Cylindrocladium and Thielaviopsis rots on woody ornamentals such as azaleas, rhododendrons, conifers, and poinsettias. Use 1 lb per 100 gals, apply as a drench or heavy spray (1 to 2 pts per sq ft) after transplanting into propagation beds or containers. Repeat at 2- to 4-week intervals during periods favorable for disease. "Tersan" 1991 DF does not control *Pythium* spp. or *Phytophthora* spp.

Preplant Dip Treatment—For diseases listed under Drench Treatment, use 1 lb per 100 gals of water, immerse plants or cuttings for 10 to 15 min, remove and allow to drain.

BULBS (Easter Lily, Tulip, Gladiolus, Daffodil, Iris): Fusarium and Penicillium Rots—Use 12.3 lbs per 100 gals of water (12 tablespoons per gal). Soak cleaned bulbs for 15 to 30 minutes in warm dip (80 to 85 F), preferably within 48 hours after digging. Dry bulbs after treatment. If bulbs are for forcing, treat after bulbs have been heat cured.

Registered trademark of Merck & Co.
 Registered trademark of Nippon Soda Co. Japan
 Registered trademark of Malsin, Inc.
 Registered trademark of DuPont Turfmark Co.

BEST AVAILABLE COPY