



8-11-71  
17-10-71

# BENOMYL 50W

**FOR PROCESSORS AND FORMULATORS ONLY**

**ACTIVE INGREDIENT**

Benomyl [Methyl 1-(butylcarbamoyl)-2-benzimidazolecarbamate] . . .50%

**INERT INGREDIENTS** . . . . .50%

U.S. Pat. Pending

USDA Reg. No. 352-358

Keep out of reach of children.

**CAUTION!** MAY IRRITATE EYES, NOSE, THROAT, AND SKIN.

Avoid breathing dust. Avoid contact with skin, eyes, and clothing.

Wash thoroughly after using. Keep away from fire or sparks.

In case of contact, flush skin or eyes with plenty of water; for eyes, get medical attention.

**IMPORTANT** — Never allow Benomyl 50W to become wet during storage. This may lead to certain chemical changes which will reduce the effectiveness of Benomyl 50W as a fungicide. Keep container closed when not in use.

**NET 110 LBS.**

**ACCEPTED**  
  
MAY 4 - 1971  
  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTER-  
ED UNDER NO. 352-358

GC-20007 4-71

Made in U.S.A. Printed in U.S.A.

E. I. DU PONT DE NEMOURS & CO INC INDUSTRIAL AND BIOCHEMICALS DEPT WILMINGTON DE

To prevent sucker development on flue-cured and burley tobacco: MH-30<sup>®</sup>, a plant inhibitor, offers a new method of controlling sucker growth by keeping suckers from growing rather than by killing them. Since MH-30<sup>®</sup> is absorbed and moves inside the tobacco plant, spraying the whole plant is not necessary. Best results are obtained by spraying the upper 1/3 to 1/2 of the plant. It is very important to apply equal amounts of MH-30<sup>®</sup> to each plant at time of topping.

**When to spray:** Satisfactory sucker control depends on correct timing of the spray.

- a. If all plants flower at about the same time, proceed as follows: As soon as the plants are in FULL FLOWER, top, pull all suckers from early-maturing plants and spray MH-30<sup>®</sup> uniformly onto the entire field within 24 hours after topping. (See notes 1-9).
- b. If there is a wide range in the time of flowering, the following procedure should be used:
  1. Go through the field and top the few early-flowering plants.
  2. Wait until the remaining plants are in FULL FLOWER, top them, remove all suckers from the entire field, and apply MH-30<sup>®</sup> within 24 hours after topping. (See notes 1-9.) Tip leaves to be harvested should be at least 6 inches long at time of spraying.

**IMPORTANT NOTICE** — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Seller, and Buyer assumes the risk of any such use.

**SHAKE WELL BEFORE USING!**

Lot No.

5-24-76