




# METHOMYL



## COMPOSITION

### FOR REFORMULATION USES ONLY

**ACTIVE INGREDIENT: Methomyl**

**S-methyl N-[(methylcarbamoyl)oxy]thioacetimidate 90%**

**INERT INGREDIENTS 10%**

EPA REG. NO. 352-367

EPA REG. NO. 352-367 AA

Keep out of reach of children.

**DANGER**



**POISON**

**MAY BE FATAL IF SWALLOWED. POISONOUS IF INHALED.**

**Do not breathe dust.**

**Do not get in eyes, on skin, on clothing.**

In case of contact, wash skin with plenty of soap and water; for eyes, flush with water for 15 minutes and get medical attention; remove and wash contaminated clothing before re-use.

Wear goggles and a mask or respirator of a type suitable for protection against insecticide dust when handling. Wear clean clothes daily. Wash thoroughly after handling and before eating or smoking. Do not contaminate or store near feed or foodstuffs.

**IMPORTANT**—Keep unprotected persons out of operational areas. Do not re-use container; crush and bury when empty.

This product is toxic to fish, birds and other wildlife. Keep out of any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes.

## ANTIDOTE AND FIRST AID

**ATROPINE IS AN ANTIDOTE—CONSULT PHYSICIAN FOR EMERGENCY SUPPLY OF 1/100 GRAIN ATROPINE TABLETS. CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED POISONING.**

If **swallowed**, give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. If **inhaled**, remove from exposure. Have patient lie down and keep quiet. If patient is not breathing, start artificial respiration immediately. **Never give anything by mouth to an unconscious person.**

If **warning symptoms appear** (see **Warning Symptoms** below), before physician arrives, immediately swallow two atropine tablets (each 1/100 gr.); thereafter, every 10 to 15 minutes, take one atropine tablet (1/100 gr.) until throat becomes dry and skin becomes dry and flushed. Take additional tablets as necessary to maintain a moderately dry throat and dry, flushed skin until physician arrives.

## NOTE TO PHYSICIAN

**WARNING SYMPTOMS**—Methomyl poisoning produces effects associated with anticholinesterase activity which may include weakness, blurred vision, headache, nausea, abdominal cramps, discomfort in the chest, constriction of pupils, sweating, slow pulse, muscle tremors.

**TREATMENT**—Atropine sulfate should be used for treatment. Administer repeated doses, 1.2 to 2.0 mg. intravenously every 10 to 30 minutes until full atropinization is achieved. Maintain atropinization until the patient recovers. Artificial respiration or oxygen may be necessary. Allow no further exposure to any cholinesterase inhibitor until recovery is assured. Do not use morphine or 2-PAM.

**NET LBS.**

E. I. DU PONT DE NEMOURS & CO. (INC.), BIOCHEMICALS DEPARTMENT, WILMINGTON, DELAWARE