



IMPORTANT: Wear bandana, wash goggles and mask when using this product. For protection against dust, use dust mask. Wash face before and after use. Use 11 P. Cartridge or Mini Safety Applicator. Do not use in enclosed spaces. Wash thoroughly after handling. Do not eat, drink or smoke while using. Store in a cool, dry place.

Keep in protected place. Do not permit children to handle. Do not use in or near water. Bury when empty.

This product is toxic to fish, birds and other wildlife. Keep out of reach of children. Do not contaminate water by cleaning of equipment or disposal of waste.

ACCEPTED
NOV 07 1968
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
Reg. No. 352-361

8-21561 3-77

Made in U.S.A.

Printed in U.S.A.



METHOMYL COMPOSITION

For Formulation of Insecticides Only

ACTIVE INGREDIENT: Methomyl
S-methyl N[(methylcarbamoyl)oxy]thioacetimidate 90%
INERT INGREDIENTS 10%
U. S. Pat. 3,576,824 & 3,639,633 EPA Reg. No. 352-361-AA
EPA Est. 352-TX-1

Keep out of reach of children.

DANGER  **POISON**

See Antidote in Right Panel ▶

MAY BE FATAL IF SWALLOWED. POISONOUS IF INHALED.

Do not breathe dust.

Do not get in eyes, on skin, on clothing.

In case of contact, wash skin with plenty of soap and water, for eyes, flush with water for 15 minutes and get medical attention, remove and wash contaminated clothing before re use.

NET 2 LBS.

5 E. I. du Pont de Nemours & Co. (Inc.)
BIOCHEMICALS DEPARTMENT, WILMINGTON, DELAWARE

ATRO:
GRAIN
POISON

If swallowed
throat
to an

If inhaled
breath
UNCOM

If worn
immed
minute
dry an
and d

WARN
estera
inal cr
tremo

TREAT
1.2 to
Mainta
be nec
assur

Do no
phoro
treatm

NOTIC
of con



METHOMYL COMPOSITION

For Formulation of Insecticides Only

ACTIVE INGREDIENT: Methomyl
S-methyl N-[(methylcarbamoyl)oxy]thioacetimidate 90%
INERT INGREDIENTS 10%
U. S. Pats. 3,576,834 & 3,639,633 EPA Reg. No. 352-361-AA
EPA Est. 352-TX-1

Keep out of reach of children.

DANGER  **POISON**

See Antidote in Right Panel ▶

MAY BE FATAL IF SWALLOWED. POISONOUS IF INHALED.

Do not breathe dust.

Do not get in eyes, on skin, or on clothing.

In case of contact, wash skin with plenty of soap and water; for eyes, flush with water for 15 minutes and get medical attention; remove and wash contaminated clothing before re-use.

NET 2 LBS.

5 E. I. du Pont de Nemours & Co. (Inc.)
BIOCHEMICALS DEPARTMENT, WILMINGTON, DELAWARE



POISON



ANTIDOTE AND FIRST AID

ATROPINE IS AN ANTIDOTE—CONSULT PHYSICIAN FOR EMERGENCY SUPPLY OF 1/100 GRAIN ATROPINE TABLETS. CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED POISONING.

If swallowed, drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger or blunt object. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

If inhaled, remove from exposure. Have patient lie down and keep quiet. If patient is not breathing, start artificial respiration immediately. Never give anything by mouth to an unconscious person.

If warning symptoms appear (See warning symptoms below), before physician arrives, immediately swallow two atropine tablets (each 1/100 gr.); thereafter, every 10 to 15 minutes, take one atropine tablet (1/100 gr.) until throat becomes dry and skin becomes dry and flushed. Take additional tablets as necessary to maintain a moderately dry throat and dry, flushed skin until physician arrives.

NOTE TO PHYSICIAN

WARNING SYMPTOMS—Methomyl poisoning produces effects associated with anticholinesterase activity which may include weakness, blurred vision, headache, nausea, abdominal cramps, discomfort in the chest, constriction of pupils, sweating, slow pulse, muscle tremors.

TREATMENT—Atropine sulfate should be used for treatment. Administer repeated doses, 1.2 to 2.0 mg. intravenously every 10 to 30 minutes until full atropinization is achieved. Maintain atropinization until the patient recovers. Artificial respiration or oxygen may be necessary. Allow no further exposure to any cholinesterase inhibitor until recovery is assured.

Do not use morphine or 2-PAM. However, for exposure to Methomyl and organophosphorous insecticides, 2-PAM may be used as required to supplement the atropine sulfate treatment as described above.

NOTICE TO BUYER—Purchase of this material does not confer any rights under patents of countries outside of the United States.