

REPRODUCTION LABEL ACTUAL SIZE

BEST DOCUMENT AVAILABLE

REPRODUCTION

NOTICE OF WARRANTY

Du Pont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. On injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, any of which are beyond the control of Du Pont. In no case shall Du Pont be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. DU PONT MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

DIRECTIONS

Du Pont "Lannate" 4-G Methomyl Insecticide Granular should be used only in accordance with recommendations on this label, or in separate published Du Pont recommendations available through local dealers.

Du Pont "Lannate" 4-G Methomyl Insecticide Granular is designed for control of the south western corn borer in sorghum in the State of Arizona. The granules may be applied in ground or aerial equipment to commercial plantings only. Do not use in home plantings.

Application

Make a broadcast application at the rate of 10 to 25 lbs. per acre when approximately 75% of the whorls are infested with worms but before the worms bore into the stalk of the sorghum. A second treatment may be applied if needed but the total amount applied should not exceed 25 lbs. per acre. Do not apply after sorghum has reached the boot stage. Do not feed, forage or ensile immature sorghum.

IMPORTANT



Lannate® 4-G METHOMYL INSECTICIDE GRANULAR

ACTIVE INGREDIENT: Methomyl S-methyl N-[(methylcarbamoyl)oxy]thioacetimidate 4% INERT INGREDIENTS 96% U.S. Pat. 2,576,834 & 2,639,633 EPA Reg. No. 352-363 EPA Est. 352-WV-1

Keep out of reach of children.

WARNING! HARMFUL IF SWALLOWED OR INHALED.

Do not breathe dust. Do not get in eyes, on skin, or clothing.

In case of contact, wash skin with plenty of soap and water; for eyes, flush with water for 15 minutes and get medical attention; remove and wash contaminated clothing before re use.

NET 10 LBS.

E. I. DU PONT DE NEMOURS & COMPANY (INC.) BIOCHEMICALS DEPARTMENT WILMINGTON, DELAWARE

ANTIDOTE AND

ATROPINE IS AN ANTIDOTE. CONSULT PHYSICIAN FOR GRAIN ATROPINE TABLETS. CALL A PHYSICIAN IN CASE OF POISONING.

If swallowed, give a tablespoonful of salt water. If fluid is clear. If inhaled, remove from exposure. If patient is not breathing, start artificial respiration. If mouth to an unconscious person.

If warning symptoms appear (See warning symptoms) immediately swallow two atropine tablets (each containing 1/100 gr.) until the patient is flushed. Take additional tablets as necessary until flushed skin until physician arrives.

NOTE TO

Warning Symptoms—Methomyl poisoning produces activity which may include weakness, blurred vision, discomfort in the chest, constriction of pupils.

Treatment—Atropine sulfate should be used for 20 mg. intravenously every 10 to 30 minutes until atropinization until the patient recovers. Allow no further exposure to any cholinesterase inhibitors.

Do not use in

NOTICE TO BUYER: Purchase of this material is restricted to the United States.

B 21420 12 75

BEST DOCUMENT AVAILABLE

ACTUAL SIZE

BEST DOCUMENT AVAILABLE



Lannate 4-G

METHOMYL INSECTICIDE
GRANULAR

ACTIVE INGREDIENT: Methomyl
S-methyl N-[(methylcarbamoyl)oxy]thioacetimdate 4%
INERT INGREDIENTS 96%
U.S. Pat. 2,576,834 & 2,637,433 EPA Reg. No. 352-363
EPA Est. 352-WV-1

Keep out of reach of children.

WARNING! HARMFUL IF SWALLOWED OR INHALED.

Do not breathe dust
Do not get in eyes, on skin, on clothing

In case of contact, wash skin with plenty of soap and water; for eyes, flush with water for 15 minutes and get medical attention; remove and wash contaminated clothing before re use

NET 10 LBS.

E. I. DU PONT DE NEMOURS & COMPANY (INC.)
BIOCHEMICALS DEPARTMENT
WILMINGTON, DELAWARE

17

W

BEST DOCUMENT AVAILABLE

ACTUAL SIZE

ANTIDOTE AND FIRST AID

ATROPINE IS AN ANTIDOTE. CONSULT PHYSICIAN FOR EMERGENCY SUPPLY OF 1/100 GRAIN ATROPINE TABLETS. CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED POISONING.

If swallowed, give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. If inhaled, remove from exposure. Have patient lie down and keep quiet. If patient is not breathing, start artificial respiration immediately. Never give anything by mouth to an unconscious person.

If warning symptoms appear (See warning symptoms below), before physician arrives, immediately swallow two atropine tablets (each 1/100 gr.); thereafter, every 10 to 15 minutes, take one atropine tablet (1/100 gr.) until throat becomes dry and skin becomes dry and flushed. Take additional tablets as necessary to maintain a moderately dry throat and dry, flushed skin until physician arrives.

NOTE TO PHYSICIAN

Warning, Symptoms—Methomyl poisoning produces effects associated with anticholinesterase activity which may include weakness, blurred vision, headache, nausea, abdominal cramps, discomfort in the chest, constriction of pupils, sweating, slow pulse, muscle tremors

Treatment—Atropine sulfate should be used for treatment. Administer repeated doses, 1.2 to 2.0 mg. intravenously every 10 to 30 minutes until full atropinization is achieved. Maintain atropinization until the patient recovers. Artificial respiration or oxygen may be necessary. Allow no further exposure to any cholinesterase inhibitor until recovery is assured.

Do not use morphine or 2 PAM

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.

00 18

B 21420 127

Printed in U.S.A.

BEST DOCUMENT AVAILABLE