

SPECIMEN LABEL ACTUAL SIZE

NOTICE OF WARRANTY

Du Pont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Du Pont. In no case shall Du Pont be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. DU PONT MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

DIRECTIONS

Du Pont "Lannate" G Methomyl Insecticide Granular should be used only in accordance with recommendations on this label, or in separate published Du Pont recommendations available through local dealers.

"Lannate" G is a granular product for control of the southwestern corn borer in sorghum in the State of Arizona. "Lannate" G may be applied in ground or aerial equipment to commercial plantings only. Do not use in home plantings.

Application

Apply 8 to 20 lbs. per acre when approximately 25% of the whorls are infested with worms but before the worms bore into the stalk of the sorghum. A second treatment may be applied if needed but the total amount applied should not exceed 20 lbs. per acre. Do not apply after the sorghum has reached the boot stage. Do not feed, forage or ensile immature sorghum.

IMPORTANT

BEST DOCUMENT AVAILABLE



Lannate G METHOMYL INSECTICIDE GRANULAR


ACTIVE INGREDIENT: Methomyl S-methyl N-[(methylcarbamoyl)oxy]thioacetimidate ... 5% INERT INGREDIENTS ... 95% U.S. Pat. 3,576,834 & 3,639,633 EPA Reg. No. 352-364 EPA Est. 352-WV-1

Keep out of reach of children.

WARNING! HARMFUL IF SWALLOWED OR INHALED.

Use as directed. Do not use in home plantings.

In case of contact, wash skin with plenty of soap and water; for eyes, flush with water for 15 minutes and get medical attention; remove and wash contaminated clothing before re-use.



NET 10 LBS.

E. I. DU PONT DE NEMOURS & COMPANY (INC.) BIOCHEMICALS DEPARTMENT WILMINGTON, DELAWARE

NOTICE TO BUYER: Purchase of this material outside of the United States

8 21453 4-74

SPECIMEN

ANTIDOTE AT

ATROPINE IS AN ANTIDOTE CONSULT PHYSICIAN. GRAB ATROPINE TABLETS. CALL A PHYSICIAN. POISONING.

If swallowed, give a tablespoonful of salt in fluid is clear. If inhaled, remove from exposure. If patient is not breathing, start artificial respiration. If unconscious, flush mouth to an unconscious person.

If warning symptoms appear (See warning symptoms) immediately swallow two atropine tablets (each 1 mg.) take one atropine tablet (1/100 gr.) until flushed. Take additional tablets as necessary. Flush skin until physician arrives.

NOTE TO

Warning Symptoms--Methomyl poisoning produces activity which may include weakness, blurred vision, discomfort in the chest, constriction of pupils.

Treatment--Atropine sulfate should be used 2.0 mg. intravenously every 10 to 30 minutes until the patient recovers. After recovery allow no further exposure to any cholinesterase.

Do not use more

BEST DOCUMENT AVAILABLE

LABEL ACTUAL SIZE

WARRANTY

Read the chemical description on the label stated on such label only when used in the conditions. It is impossible to eliminate of this product. Crop injury, inefficiency may result because of such factors as soil, or the manner of use or application, but in no case shall Du Pont be liable as resulting from the use or handling of it by the Buyer. DU PONT MAKES NO CLAIMS FOR A PARTICULAR PURPOSE NOR FOR EXCEPT AS STATED ABOVE

INSTRUCTIONS

Granular should be used only in accordance with separate published Du Pont recommendations.

Use on the southwestern corn borer in sorghum in ground or aerial equipment in commercial quantities.

25% of the whorls are infested with worms in sorghum. A second treatment may be applied if exceed 20 lbs. per acre. Do not apply after harvest, forage or ensile immature sorghum.

CAUTION

BEST DOCUMENT AVAILABLE



Lannate® G

METHOMYL INSECTICIDE
GRANULAR

ACTIVE INGREDIENT: Methomyl
S-methyl N-[(methylcarbamoyl)oxy]thioacetimidate . . . 5%
INERT INGREDIENTS 95%
U.S. Pat. 3,574,834 & 3,639,633 EPA Reg. No. 352-344
EPA Est. 352-WV 1

Keep out of reach of children.

WARNING! HARMFUL IF SWALLOWED OR INHALED.

In case of contact, wash skin with plenty of soap and water; for eyes, flush with water for 15 minutes and get medical attention; remove and wash contaminated clothing before re use.

NET 10 LBS.

E. I. DU PONT DE NEMOURS & COMPANY (INC.)
BIOCHEMICALS DEPARTMENT
WILMINGTON, DELAWARE

AC
40

SPECIMEN LABEL ACTUAL SIZE

ANTIDOTE AND FIRST AID

ATROPINE IS AN ANTIDOTE CONSULT PHYSICIAN FOR EMERGENCY SUPPLY OF 1/100 GRAIN ATROPINE TABLETS CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED POISONING

If swallowed, give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. If inhaled, remove from exposure. Have patient lie down and keep quiet. If patient is not breathing, start artificial respiration immediately. Never give anything by mouth to an unconscious person.

If warning symptoms appear (See warning symptoms below), before physician arrives, immediately swallow two atropine tablets (each 1/100 gr.); thereafter, every 10 to 15 minutes, take one atropine tablet (1/100 gr.) until throat becomes dry and skin becomes dry and flushed. Take additional tablets as necessary to maintain a moderately dry throat and dry, flushed skin until physician arrives.

NOTE TO PHYSICIAN

Warning Symptoms—Methomyl poisoning produces effects associated with anticholinesterase activity which may include weakness, blurred vision, headache, nausea, abdominal cramps, discomfort in the chest, constriction of pupils, sweating, slow pulse, muscle tremors.

Treatment—Atropine sulfate should be used for treatment. Administer repeated doses, 1.2 to 2.0 mg. intravenously every 10 to 30 minutes until full atropinization is achieved. Maintain atropinization until the patient recovers. Artificial respiration or oxygen may be necessary. Allow no further exposure to any cholinesterase inhibitor until recovery is assured.

Do not use morphine or 2 PAM

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States

41
Printed in U.S.A.

BEST DOCUMENT AVAILABLE