




REG. U.S. PAT. OFF.

# METHOMYL

## LIQUID CONCENTRATE

### FOR REFORMULATION USES ONLY



**ACTIVE INGREDIENT: Methomyl**

**S-methyl N-[(methylcarbamoyl)oxy]thioacetimidate**

26%

**INERT INGREDIENTS**

74%

Keep out of reach of children.

**DANGER**



**POISON**

**FLAMMABLE**

**MAY BE FATAL IF SWALLOWED OR INHALED**

**CAUSES EYE IRRITATION**

**Do not get in eyes, on skin, on clothing.**

**Do not breathe mist or vapors.**

**Keep away from heat, sparks and open flame.**

**Keep container closed. Use with adequate ventilation.**

In case of contact, wash skin with plenty of soap and water. For eyes, flush with water for 15 minutes and get medical attention; remove and wash contaminated clothing before reuse.

Wear goggles and a mask or respirator of a type suitable for protection against insecticide mist or vapors when handling. Wear clean clothes daily. Wash thoroughly after handling and before eating or smoking. Do not contaminate water or store near food or foodstuffs.

**IMPORTANT** Keep unprotected persons out of operation areas.

This product is toxic to fish, birds and other wildlife. Keep out of any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes.

### ANTIDOTE AND FIRST AID

**ATROPINE IS AN ANTIDOTE—CONSULT PHYSICIAN FOR EMERGENCY SUPPLY OF 1-100 GRAIN ATROPINE TABLETS. CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED POISONING.**

If swallowed, give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. If inhaled, remove from exposure. Have patient lie down and keep quiet. If patient is not breathing, start artificial respiration immediately. **Never give anything by mouth to an unconscious person.**

If warning symptoms appear (see **Warning Symptoms** below), before physician arrives, immediately swallow two atropine tablets (each 1-100 gr.), thereafter, every 10 to 15 minutes, take one atropine tablet (1-100 gr.) until throat becomes dry and skin becomes dry and flushed. Take additional tablets as necessary to maintain a moderately dry throat and dry, flushed skin until physician arrives.

### NOTE TO PHYSICIAN

**WARNING SYMPTOMS** Methomyl poisoning produces effects associated with anticholinesterase activity which may include weakness, blurred vision, headache, nausea, abdominal cramps, diarrhea, ataxic gait, incontinence of urine, sweating, slow pulse, muscle tremors.

**TREATMENT** Atropine sulfate should be used for treatment. Administer repeated doses, 1-2 to 2.0 mg. intravenously every 10 to 30 minutes until full atropinization is achieved. Maintain atropinization until the patient recovers. Artificial respiration or oxygen may be necessary. Allow no further exposure to any cholinesterase inhibitor until recovery is assured. Do not use morphine or 2-PAM.

**NET**

**GALS.**


E. I. DU PONT DE NEMOURS & CO. (INC.), BIOCHEMICALS DEPARTMENT, WILMINGTON, DELAWARE



# METHOMYL

## LIQUID CONCENTRATE

FOR REFORMULATION USES ONLY



ACTIVE INGREDIENT: Methomyl	26%
S-methyl N-[(methylcarbamoyl)oxy]thioacetimidate	
INERT INGREDIENTS	74%

Keep out of reach of children.

**DANGER**  **POISON**  
**FLAMMABLE**  
**MAY BE FATAL IF SWALLOWED OR INHALED**  
**CAUSES EYE IRRITATION**

Do not get in eyes, on skin, on clothing.  
Do not breathe mist or vapors.  
Keep away from heat, sparks and open flame.  
Keep container closed. Use with adequate ventilation.

Use only for agricultural purposes. Do not use for household purposes. Do not use on food crops. Do not use on ornamental plants. Do not use on lawns. Do not use on trees. Do not use on shrubs. Do not use on flowers. Do not use on vegetables. Do not use on fruits. Do not use on nuts. Do not use on seeds. Do not use on grain. Do not use on hay. Do not use on straw. Do not use on manure. Do not use on compost. Do not use on mulch. Do not use on soil. Do not use on water. Do not use on air. Do not use on earth. Do not use on sky. Do not use on sun. Do not use on moon. Do not use on stars. Do not use on planets. Do not use on galaxies. Do not use on the universe.

When using, wear protective clothing and equipment. Avoid contact with skin, eyes, and clothing. Do not breathe mist or vapors. Do not get on skin, on clothing, or in eyes. Do not get in eyes, on skin, on clothing. Do not breathe mist or vapors. Do not get on skin, on clothing, or in eyes. Do not get in eyes, on skin, on clothing. Do not breathe mist or vapors. Do not get on skin, on clothing, or in eyes.

**IMPORTANT:** Keep out of reach of children. Do not use for household purposes. Do not use on food crops. Do not use on ornamental plants. Do not use on lawns. Do not use on trees. Do not use on shrubs. Do not use on flowers. Do not use on vegetables. Do not use on fruits. Do not use on nuts. Do not use on seeds. Do not use on grain. Do not use on hay. Do not use on straw. Do not use on manure. Do not use on compost. Do not use on mulch. Do not use on soil. Do not use on water. Do not use on air. Do not use on earth. Do not use on sky. Do not use on sun. Do not use on moon. Do not use on stars. Do not use on planets. Do not use on galaxies. Do not use on the universe.

BEST DOCUMENT AVAILABLE NET GALS.

E. I. DU PONT DE NEMOURS & CO. (INC.), BIOCHEMICALS DEPARTMENT, WILMINGTON, DELAWARE

### ANTIDOTE AND FIRST AID

ATROPINE IS AN ANTIDOTE—CONSULT PHYSICIAN FOR EMERGENCY SUPPLY OF 1-100 GRAIN ATROPINE TABLETS. CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED POISONING.

If swallowed, induce vomiting with 1 cup of water. Do not repeat until 15 minutes after the first vomiting. If inhaled, remove the person to fresh air. If necessary, give artificial respiration. Never give anything by mouth to an unconscious person.

If warning symptoms appear, Warning Symptoms: Irritation of eyes, nose, throat, and skin. Headache, dizziness, blurred vision, dry mouth, and difficulty in swallowing. If these symptoms appear, stop using immediately and seek medical attention.

### NOTE TO PHYSICIAN

**WARNING SYMPTOMS:** Irritation of eyes, nose, throat, and skin. Headache, dizziness, blurred vision, dry mouth, and difficulty in swallowing. If these symptoms appear, stop using immediately and seek medical attention.

**TREATMENT:** If swallowed, induce vomiting with 1 cup of water. Do not repeat until 15 minutes after the first vomiting. If inhaled, remove the person to fresh air. If necessary, give artificial respiration. Never give anything by mouth to an unconscious person.