

## **PLEASE NOTE**

**This image contains more than one label approved for this product on this date.**

352-370

03/11/1975



1/1



# AGRICULTURAL BULLETIN

LANNATE® L METHOMYL INSECTICIDE  
SORGHUM

ACCEPTED

MAR 11 1975

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 352-370

SUPPLEMENTAL LABELING  
EPA REG. NO. 352-370

LANNATE® L METHOMYL INSECTICIDE  
FOR USE ON SORGHUM



Du Pont "Lannate" L Methomyl Insecticide is recommended for use on sorghum for control of corn earworm.

### HOW TO USE

Apply 1 to 2 pints "Lannate" L per acre in sufficient water (5 to 15 gals. per acre by air) to obtain uniform coverage. Apply when worms appear in damaging numbers and repeat at 5- to 7-day intervals or as needed.

NOTE: Do not apply within 14 days of harvest. Do not graze or feed treated plants to livestock within 14 days after application.

### IMPORTANT

BEFORE USING "LANNATE" L, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

12/12/74

*data submitted 12/19/74; accepted 3/11/75 RSV+POH*

**NEXT**

**LABEL**

352-370

3-11-1975



# AGRICULTURAL BULLETIN



LANNATE® L METHOMYL INSECTICIDE  
CARROTS  
ARIZONA AND CALIFORNIA

ACC  
MAR 11 1975  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 352-370

SUPPLEMENTAL LABELING  
EPA REG. NO. 352-370



LANNATE® L METHOMYL INSECTICIDE  
FOR USE ON CARROTS  
IN THE STATES OF ARIZONA AND CALIFORNIA

Du Pont "Lannate" L Methomyl Insecticide is recommended for use on carrots to control beet armyworm in the states of Arizona and California.

### HOW TO USE

Apply 1 1/3 to 2 pints "Lannate" L per acre in sufficient water (5 to 15 gals. per acre by air) to obtain uniform coverage. Make first application when insects first appear and repeat at 5- to 7-day intervals or as needed.

NOTE: Do not apply within 1 day of harvest.

### IMPORTANT

BEFORE USING "LANNATE" L, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

12/12/74

*data submitted 12/19/74; accepted 3/11/75 POH*

*0.2 g per root vegetable E.P. 2/26/74*


**NEXT**

**LABEL**

352-370

03/11/1975

1/1



# AGRICULTURAL BULLETIN



LANNATE® L METHOMYL INSECTICI  
APPLES  
EAST OF ROCKY MOUNTAINS

**ACCEPTED**  
MAR 11 1975  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 352-370

SUPPLEMENTAL LABELING  
EPA REG. NO. 352-370

LANNATE® L METHOMYL INSECTICIDE  
FOR USE ON APPLES  
IN AREAS EAST OF THE ROCKY MOUNTAINS



Du Pont "Lannate" L Methomyl Insecticide is recommended for use on apples to control certain aphids, codling moth, and red-banded leafroller in areas east of the Rocky Mountains.

### HOW TO USE

Apple Aphid, Rosy Apple Aphid - Apply 1 to 2 pints per 100 gals. to established infestations.

Codling Moth - Make 4 to 7 applications of 1 to 2 pints per 100 gals. at 10- to 12-day intervals beginning at the second cover spray.

Red-Banded Leafroller (Northeast and Central U.S. Only) - Apply 2 pints per 100 gals. at green tip to 1/2 inch green and at petal fall and first cover.

Apply in sufficient water to obtain thorough coverage (run-off) but not in excess of 400 gals. per acre. For concentrate sprayers, use proportionately less water.

Do not apply within 28 days of harvest. Do not graze livestock in treated orchards for 10 days after treatment.

### IMPORTANT


BEFORE USING "LANNATE" L, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

*data submitted 12/19/74 by inference from 352-372 acceptance; accepted 3/11/75 F011*

**NEXT**

**LABEL**



352-370


03/11/1975  
DUPONT

# AGRICULTURAL BULLETIN

LANNATE® L METHOMYL INSECTICIDE  
TURF - STATE OF FLORIDA

**ACCEPTED**  
**MAR 11 1975**  
Under the Federal Insecticide, Fungicide, and Plant Disease Act, as amended, for use on turf registered under EPA Reg. No. 352-370

SUPPLEMENTAL LABELING  
EPA REG. NO. 352-370



LANNATE® L METHOMYL INSECTICIDE  
FOR USE ON TURF  
IN THE STATE OF FLORIDA

Du Pont "Lannate" L Methomyl Insecticide is recommended for use on turf for control of sod webworm in the State of Florida.

### HOW TO USE

Apply 4 to 8 pints per acre (1.5 to 3.0 fluid ounces per 1,000 square feet). After application, sprinkler irrigate for 15 minutes to wash the "Lannate" L into the turf. Apply when worms first appear and repeat as needed.

NOTE: Not for use in or around the home. Do not graze or feed clippings from treated areas to livestock.

### IMPORTANT

BEFORE USING "LANNATE" L, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

12/12/74

12/19/74 accepted POH