

PLEASE NOTE

This image contains more than one label approved for this product on this date.

352-370

07/15/1976




1/1



AGRICULTURAL BULLETIN

LANNATE® L
METHOMYL INSECTICIDE
BEANS
CALIFORNIA

SUPPLEMENTAL LABELING
EPA REG. NO. 352-370



LANNATE® L METHOMYL INSECTICIDE
FOR USE ON BEANS
IN CALIFORNIA

Du Pont "Lannate" L Methomyl Insecticide is recommended for use on beans for control of lygus bugs in California.

HOW TO USE

Apply 2 to 4 pints "Lannate" L per acre in sufficient water (5 to 15 gals. per acre by air) to obtain thorough, uniform coverage. Make applications when insect first appears and repeat at 5- to 7-day intervals or as needed.

Note: Pilot should not assist in the mixing and loading operation. Vines may not be foraged within 3 days of application or cut for hay within 7 days of application. Succulent beans may be harvested 3 days after last application. Do not harvest dry beans within 25 days of last application.

IMPORTANT

BEFORE USING "LANNATE" L, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

4/7/76-A.

NEXT


LABEL



AGRICULTURAL BULLETIN



LANNATE® L
METHOMYL INSECTICIDE
COTTON
EAST OF ROCKY MOUNTAINS



SUPPLEMENTAL LABELING
EPA REG. NO. 352-370

LANNATE® L METHOMYL INSECTICIDE
FOR USE ON COTTON
EAST OF THE ROCKY MOUNTAINS

Du Pont "Lannate" L Methomyl Insecticide is recommended for occasional use with other insecticides on cotton east of the Rocky Mountains for control of cotton leafperforator and beet armyworm.

HOW TO USE

Apply 2 pints "Lannate" L per acre. Use as an occasional treatment in a regular spray schedule. Do not use more often than every 10 days. Apply in sufficient water (minimum 3 gals. per acre by air) to obtain thorough coverage.

Alternately, after the desired crop (boll load) is set on the plant, up to three treatments of 2 to 3 pints "Lannate" L per acre may be applied at 3 to 5 day intervals as needed for control of these insects.

Note: Use of "Lannate" L may redden cotton foliage. If reddening is excessive, discontinue use of "Lannate" L or alternate with insecticides other than methomyl. "Lannate" L may be applied up to 15 days before harvest. Do not graze or feed treated cotton to livestock.

IMPORTANT

BEFORE USING "LANNATE" L, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

3/24/76-A