


352-370

07/27/1976



1/2




AGRICULTURAL BULLETIN

LANNATE® L
METHOMYL INSECTICIDE
APPLES
EAST OF ROCKY MOUNTAINS

SUPPLEMENTAL LABELING
EPA REG. NO. 352-370

LANNATE® L METHOMYL INSECTICIDE
FOR USE ON APPLES
IN AREAS EAST OF THE ROCKY MOUNTAINS



Du Pont "Lannate" L Methomyl Insecticide is recommended for use on apples to control aphids, codling moth, leafrollers, tufted apple budmoth, lesser appleworm, white apple leafhopper and tentiform leafminer in areas east of the Rocky Mountains.

HOW TO USE

Apple Aphid, Rosy Apple Aphid - Apply 1 to 2 pts. per 100 gals. to established infestations and repeat as needed.


Codling Moth - Make applications of 1 to 2 pts. per 100 gals. at 10- to 12-day intervals beginning when infestations first appear.

Tufted Apple Budmoth - Apply 1 to 2 pts. per 100 gals. when insects first appear and repeat treatment as needed to maintain control.

Fruittree Leafroller, Obliquebanded Leafroller, Redbanded Leafroller, Variegated Leafroller, Lesser Appleworm, White Apple Leafhopper, Tentiform Leafminer - Apply 2 pts. per 100 gals. when insects first appear and repeat treatments as needed to maintain control.

Apply in sufficient water to obtain thorough coverage (run-off) but not in excess of 400 gals. per acre. For concentrate sprayers, use proportionately less water.

Note: Do not apply to Early MacIntosh or Wealthy apple varieties. Do not apply within 8 days of harvest. Do not graze livestock in treated orchards for 10 days after treatment.



2/2

-2-

IMPORTANT

BEFORE USING "LANNATE" L, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

7/19/76

000000

000000