



# AGRICULTURAL BULLETIN



LANNATE® L METHOMYL INSECTICIDE  
COTTON

SUPPLEMENTAL LABELING  
EPA REG. NO. 352-370

LANNATE® L METHOMYL INSECTICIDE  
IN COMBINATION WITH METHYL PARATHION  
FOR USE ON COTTON  
IN THE MIDSOUTH AND SOUTHEAST

ACCEPTED  
AUG 11 1977  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under EPA Reg. No. 352-370

Du Pont "Lannate" L Methomyl Insecticide is recommended in combination with methyl parathion for use on cotton for control of bollworm and bollweevil in the Midsouth and Southeast.

## HOW TO USE

Apply 1 to 1 1/4 pts. "Lannate" L plus 1 - 2 pts. methyl parathion (4 lb./gal. emulsifiable concentrate) in sufficient water (minimum of 2 gals. per acre by air), to obtain thorough coverage. Repeat at 5- to 7-day intervals.

Apply the higher use rates and shorten spray intervals with increasing insect populations.

Note: Use of "Lannate" L in combination with methyl parathion may redden cotton foliage. If reddening is excessive, discontinue use of this combination or alternate with other insecticides. "Lannate" L + methyl parathion may be applied up to 15 days before harvest. Do not graze or feed treated cotton to livestock. Do not permit workers to handle treated crops within 48 hours after application. Do not hand pick within 7 days of application.

## IMPORTANT

BEFORE USING "LANNATE" L, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.