

PLEASE NOTE

This image contains more than one label approved for this product on this date.

352-370

11/07/1977

1/1



AGRICULTURAL BULLETIN



LANNATE® L METHOMYL INSECTICIDE
SUGAR BEETS
ARIZONA

SUPPLEMENTAL LABELING
EPA REG. NO. 352-370

ACCEPTED
NOV 7 1977
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
this product is approved for the pesticides
required under EPA Reg. No. 352-370

LANNATE® L METHOMYL INSECTICIDE
FOR USE ON SUGAR BEETS
IN ARIZONA

Du Pont "Lannate" L Methomyl Insecticide is recommended for use on sugar beets for the control of beet armyworm, beet webworm, flea beetle and aphids in Arizona.

HOW TO USE

Apply 1 to 4 pts. "Lannate" L per acre in sufficient water (minimum 3 gals. per acre by air) to obtain thorough coverage. Make the first application when insects appear and plant damage occurs. Make 2 or 3 additional applications if needed to maintain control.

For effective insect control, proper spray timing and coverage are essential. Best control is obtained when spray schedules are initiated on small insects. In general, use the low rate on light insect infestations. In beet armyworm control, use the dosage rates below 2 pts. "Lannate" L per acre only when treating small plants.


Note: Do not apply within 7 days of harvest. Do not feed treated crops to livestock within 30 days of last application.

IMPORTANT

BEFORE USING "LANNATE" L, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

7/11/77



NEXT

LABEL

352-370

11-7-1977

1/1-



AGRICULTURAL BULLETIN



LANNATE® L METHOMYL INSECTICIDE
SPINACH - FLORIDA

SUPPLEMENTAL LABELING
EPA REG. NO. 352-370

LANNATE® L METHOMYL INSECTICIDE
FOR USE ON SPINACH
IN FLORIDA

ACCEPTED
NOV 7 1977
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No. 352-370

Du Pont "Lannate" L Methomyl Insecticide is recommended for use on spinach for control of cabbage looper and beet armyworm in Florida.

HOW TO USE

Apply 2 to 4 pts. "Lannate" L per acre in sufficient water (minimum 3 gals. per acre by air) to obtain thorough coverage. Apply at 5- to 7-day intervals or as needed. Apply the lower rate on small plants, small worms and light infestations of insects. Use 1 to 3 applications of the highest recommended rate for controlling severe infestations. Thereafter, use the lowest rate possible to maintain control.

Note: Do not apply when minimum daily temperatures are 32°F. or lower. Do not apply within 7 days of harvest. Do not apply to seedling spinach when the whorl is less than 3" in diameter or injury to the crop may result.

IMPORTANT

BEFORE USING "LANNATE" L, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

7/11/77

