

## **PLEASE NOTE**

**This image contains more than one label  
approved for this product on this date.**



# AGRICULTURAL BULLETIN

**DU PONT**  
LANNATE® L INSECTICIDE  
COTTON  
EAST OF ROCKY MOUNTAINS

352-370

05/16/1978  
RESTRICTED USE PESTICIDE

1/2

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

**ACCEPTED**

MAY 16 1978

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, the pesticide registered under EPA Reg. No. 352-370

SUPPLEMENTAL LABELING  
EPA REG. NO. 352-370

LANNATE® L INSECTICIDE  
FOR USE ON COTTON  
EAST OF ROCKY MOUNTAINS

Du Pont "Lannate" L Insecticide is recommended for use on cotton to control bollworm and tobacco budworm east of the Rocky Mountains. Control is obtained through kill of eggs (ovicide) and larvae (larvicide).

## HOW TO USE

### Dosage Rates

<u>STAGE OF INSECT</u>	<u>PTS/ACRE</u>	<u>SCHEDULE OF SPRAYS</u>	<u>SPRAY TIMING</u>
Larvae + eggs of bollworm, budworm	2	Use as an occasional spray in regular schedule but not more often than every 10 days.	Best results occur on smaller larvae including those up to 1/2 inch long.
Larvae + eggs of bollworm,	2-3	Up to 3 applications at 3-5 day intervals after desired boll- load set on plants.	Best results occur on smaller larvae including those up to 1/2 inch long. Use the higher rate on severe infestations and larger larvae.

### Spray Application

Use sufficient water to obtain thorough, uniform coverage. For aerial application, use a minimum of 3 gals. per acre.

### Use Restrictions

"Lannate" L may be applied up to 15 days before harvest. Do not graze or feed treated cotton to livestock. Use of "Lannate" L

may redden cotton foliage. If reddening is excessive, alternate with other insecticides or discontinue use.

2/2

IMPORTANT

BEFORE USING "LANNATE" L, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

9/19/77-A

**NEXT**

**LABEL**



# AGRICULTURAL BULLETIN

352-370

5-16-1978



LANNATE® L INSECTICIDE  
COTTON

## RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

ACCEPTED

MAY 16 1978

SUPPLEMENTAL LABELING  
EPA REG. NO. 352-370

LANNATE® L INSECTICIDE  
IN COMBINATION WITH METHYL PARATHION  
FOR USE ON COTTON

352-370

Du Pont "Lannate" L Insecticide is recommended in combination with methyl parathion for use on cotton for control of tobacco budworm and bollweevil. Control of budworm with "Lannate" L is obtained through kill of eggs (ovicide) and larvae (larvicide).

### HOT TO USE

Apply 1 to 1 1/4 pts. "Lannate" L plus 1 to 2 pts. methyl parathion (4 lb./gal. emulsifiable concentrate) in sufficient water (minimum of 2 gals. per acre by air), to obtain thorough coverage. Repeat at 5- to 7-day intervals.

Apply the higher use rates and shorten spray intervals with increasing insect populations.

Note: Use of "Lannate" L in combination with methyl parathion may redden cotton foliage. If reddening is excessive, discontinue use of this combination or alternate with other insecticides. "Lannate" L + methyl parathion may be applied up to 15 days before harvest. Do not graze or feed treated cotton to livestock. Do not permit workers to handle treated crops within 48 hours after application. Do not hand pick within 7 days of application.

### IMPORTANT

BEFORE USING "LANNATE" L, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

3/16/77-A

**NEXT**

**LABEL**



# AGRICULTURAL BULLETIN

352-370 5-16-1978

LANNATE® L INSECTICIDE  
COTTON 1/1

## RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

**ACCEPTED**

**MAY 16 1978**

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, this pesticide  
registered under  
EPA Reg. No. 352-370

SUPPLEMENTAL LABELING  
EPA REG. NO. 352-370

LANNATE® L INSECTICIDE  
IN COMBINATION WITH METHYL PARATHION  
FOR USE ON COTTON  
IN THE MIDSOUTH AND SOUTHEAST

Du Pont "Lannate" L Insecticide is recommended in combination with methyl parathion for use on cotton for control of bollworm and bollweevil in the Midsouth and Southeast. Control of bollworm with "Lannate" L is obtained through kill of eggs (ovicide) and larvae (larvicide).

### HOW TO USE

Apply 1 to 1 1/4 pts. "Lannate" L plus 1 to 2 pts. methyl parathion (4 lbs./gal. emulsifiable concentrate) in sufficient water (minimum of 2 gals. per acre by air), to obtain thorough coverage. Repeat at 5- to 7-day intervals.

Apply the higher use rates and shorten spray intervals with increasing insect populations.

**Note:** Use of "Lannate" L in combination with methyl parathion may redden cotton foliage. If reddening is excessive, discontinue use of this combination or alternate with other insecticides. "Lannate" L + methyl parathion may be applied up to 15 days before harvest. Do not graze or feed treated cotton to livestock. Do not permit workers to handle treated crops within 48 hours after application. Do not hand pick within 7 days of application.

### IMPORTANT

**BEFORE USING "LANNATE" L, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.**

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

6/20/77-A