


PLEASE NOTE

**This image contains more than one label
approved for this product on this date.**



AGRICULTURAL BULLETIN

LANNATE® L INSECTICIDE
LETTUCE

352-370

07/14/1978
RESTRICTED USE PESTICIDE

1/1

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those covered by the Certified Applicator's certification.

SUPPLEMENTAL LABELING
EPA REG. NO. 352-370

LANNATE® L INSECTICIDE
FOR USE ON LETTUCE

ACCEPTED
JUL 14 1978
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 352-370

Du Pont "Lannate" L Insecticide is recommended for use on lettuce for the control of aphids, thrips and corn earworm.

HOW TO USE

Apply 2 to 4 pts. "Lannate" L per acre in sufficient water (5 to 15 gals. per acre by air) to obtain thorough, uniform coverage. Make applications when insects first appear and repeat at 5- to 7-day intervals or as needed to maintain control.

Note: Do not apply within 10 days of harvest.

IMPORTANT

BEFORE USING "LANNATE" L, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

7/22/76-A

NEXT

LABEL



AGRICULTURAL BULLETIN

LANNATE® L INSECTICIDE
SOYBEANS

352-370

07/14/1978

RESTRICTED USE PESTICIDE

1/1

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

ACCEPTED
JUL 14 1978
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 352-370

SUPPLEMENTAL LABELING
EPA REG. NO. 352-370

LANNATE® L INSECTICIDE
FOR USE ON SOYBEANS

Du Pont "Lannate" L Insecticide is recommended for insect control on soybeans. Low use rates will control light to moderate infestations of Mexican bean beetle, green cloverworm and corn earworm and help conserve beneficial predators.

HOW TO USE

Apply 1/2 to 1 pint per acre before insect populations build to economic levels. Use the lower rate on light infestations. Make additional applications as needed to retain control. Use sufficient water to obtain thorough coverage (min. of 2 gals. per acre aerial).

Periodic scouting of fields is required to insure proper timing of spray.

Note: Do not apply within 14 days of bean harvest. Do not graze or feed treated soybean forage to livestock within 3 days of last application nor soybean hay within 7 days of last application.

IMPORTANT

BEFORE USING "LANNATE" L, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

'1/25/78