

352-370

02/22/1979

1/2



AGRICULTURAL BULLETIN



LANNATE® L INSECTICIDE
BROCCOLI, BRUSSELS SPROUTS,
CABBAGE AND CAULIFLOWER

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

SUPPLEMENTAL LABELING
EPA REG. NO. 352-370

LANNATE® L INSECTICIDE

IN COMBINATION WITH BACILLUS THURINGIENSIS
FOR USE ON BROCCOLI, BRUSSELS SPROUTS, CABBAGE AND CAULIFLOWER

Du Pont "Lannate" L Insecticide is recommended in combination with the bacterium, Bacillus thuringiensis^{1/} for control of loopers, imported cabbageworm and diamondback moth on broccoli, Brussels sprouts, cabbage and cauliflower.

HOW TO USE

LOOPERS

Apply 2 pts. "Lannate" L plus 1/2 lb. Bacillus thuringiensis^{1/} per acre.

IMPORTED CABBAGEWORM, DIAMONDBACK MOTH

Apply 1 to 2 pts. "Lannate" L plus 1/4 to 1/2 lb. Bacillus thuringiensis^{1/} per acre. Apply the higher use rates on severe insect infestations.

Apply when larval populations appear in significant numbers and continue at 5- to 7-day intervals or as needed. Apply in sufficient water (minimum 3 gals. per acre by air) to obtain thorough coverage.

Note: Do not apply within the following days of harvest:

Cabbage	- 1 day
Broccoli, Brussels sprouts and cauliflower	- 3 days

^{1/} Wettable powder containing 16,000 International Units of potency per mg (7.26 billion International Units per pound); see specific product labeling as federally registered.

IMPORTANT

BEFORE USING "LANNATE" L, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

5/18/78

