

PLEASE NOTE

This image contains more than one label approved for this product on this date.



352-370

3-23-1979

AGRICULTURAL BULLETIN



LANNATE® L INSECTICIDE
COTTON

4/1

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

RECEIVED
MAR 23 1979
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 352-370

SUPPLEMENTAL LABELING
EPA REG. NO. 352-370

LANNATE® L INSECTICIDE
FOR USE ON COTTON

Du Pont "Lannate" L Insecticide is recommended for use on cotton to control fall armyworm.

East of Rocky Mountains

Early Season

Apply 2 pts. "Lannate" L per acre; use as an occasional spray regularly but not more often than every 10 days.

Late Season (after desired boll load set on plants)

Apply 2 to 3 pts. "Lannate" L per acre, use up to 3 applications at 3- to 5-day intervals.

West of Rocky Mountains

Apply 2 to 3 pts. "Lannate" L per acre. Apply at 5- to 7-day intervals or as needed to maintain control.

Apply insufficient water (minimum 3 gals. per acre by air) to obtain thorough coverage.

Initiate treatment when significant populations of fall armyworm occur.

Note: "Lannate" L may be applied up to 15 days before harvest. Do not graze or feed treated cotton to livestock. Use of "Lannate" L may redden cotton foliage. If reddening is excessive, alternate with other insecticides or discontinue use.

IMPORTANT

BEFORE USING "LANNATE" L, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

3/2/79

NEXT

LABEL



352-370

3-23-1979

AGRICULTURAL BULLETIN



LANNATE® L INSECTICIDE
COTTON
ARIZONA

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

SUPPLEMENTAL LABELING
EPA REG. NO. 352-370

LANNATE® L INSECTICIDE
FOR USE ON COTTON
IN ARIZONA

ACCEPTED

MAR 23 1979

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 352-370

Du Pont Lannate® L Insecticide is recommended for control of cotton bollworm and tobacco budworm on cotton in Arizona. When Lannate® L is applied to eggs, control is from direct kill of eggs (ovicide) and newly-hatched larvae (larvicide).

HOW TO USE

Apply 1 pt. Lannate® L per acre in sufficient water (minimum 3 gals. per acre by air) to obtain thorough coverage. Initiate schedule of sprays when significant numbers of eggs are present. Continue at 5- to 7-day intervals while eggs are present and larval control is adequate. If significant numbers of larvae survive the treatment, use higher rates of Lannate® L as registered.

Periodic scouting of fields is required to insure proper timing of spray.

Note: Lannate® L may be applied up to 15 days before harvest. Do not graze or feed treated cotton to livestock. Use of Lannate® L may redden cotton foliage. If reddening is excessive, alternate with other insecticides or discontinue use.

IMPORTANT

BEFORE USING LANNATE® L, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

1/26/79

NEXT

LABEL



352-370

3-23-1979

AGRICULTURAL BULLETIN




LANNATE® L INSECTICIDE
COTTON
EAST OF ROCKY MOUNTAINS

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

SUPPLEMENTAL LABELING
EPA REG. NO. 352-370

LANNATE® L INSECTICIDE
FOR USE ON COTTON
EAST OF THE ROCKY MOUNTAINS



Du Pont Lannate® L Insecticide is recommended for control of cotton bollworm and tobacco budworm on cotton east of the Rocky Mountains. When Lannate® L is applied to eggs, control is from direct kill of eggs (ovicide) and newly-hatched larvae (larvicide). The low use rates help conserve beneficial predators.

HOW TO USE

Apply 1/2 to 1 pt. Lannate® L per acre in sufficient water (minimum 3 gals. per acre by air) to obtain thorough coverage. Initiate schedule of sprays when significant numbers of eggs are present. Continue at 5- to 7-day intervals while eggs are present and larval control is adequate. If significant numbers of larvae survive the treatment, use higher rates of Lannate® L as registered.

Periodic scouting of fields is required to insure proper timing of spray.

Note: Lannate® L may be applied up to 15 days before harvest. Do not graze or feed treated cotton to livestock. Use of Lannate® L may redden cotton foliage. If reddening is excessive, alternate with other insecticides or discontinue use.

IMPORTANT

BEFORE USING LANNATE® L, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

1/25/79